

53

22 JUL 1999


2-514170319  
WIPO OMPI

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : C07D 213/75, 261/16, 239/42 C07D 237/20, 277/82		A1	(11) International Publication Number: WO 93/10094 (43) International Publication Date: 27 May 1993 (27.05.93)
(21) International Application Number: PCT/SE92/00758 (22) International Filing Date: 4 November 1992 (04.11.92) (30) Priority data: 9103397-7 18 November 1991 (18.11.91) SE (71) Applicant (for all designated States except US): KABI PHARMACIA AB [SE/SE]; S-751 82 Uppsala (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): AGBACK, Karl, Hubert [SE/SE]; Trällbovägen 5, S-755 90 Uppsala (SE). AHRGREN, Leif [SE/SE]; Vinkelvägen 6, S-740 60 Örbyhus (SE). BERGLINDH, Thomas [SE/SE]; Ripvägen 5, S-756 53 Uppsala (SE). HARALDSSON, Martin [SE/SE]; Kragstallundsvägen 163, S-186 51 Vallentuna (SE). OLSOSON, Lars-Inge [SE/SE]; Kämpevägen 47, S-151 54 Södertälje (SE). SMEDEGARD, Göran [SE/SE]; Oslögtan 30, S-752 64 Uppsala (SE).		(74) Agents: BERGANDER, Håkan et al.; Kabi Pharmacia AB, S-751 82 Uppsala (SE). (81) Designated States: AU, BB, BG, BR, CA, CS, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, PL, RO, RU, SD, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published With international search report.	
(54) Title: NOVEL SUBSTITUTED SALICYCLIC ACIDS (57) Abstract <p>A compound of the formula Het-NR-SO<sub>2</sub>-Ph<sup>2</sup>-A-Ph<sup>1</sup> (COOH)(OH) and tautomeric form, salts, solvates, C<sub>1-6</sub> alkyl esters and pharmaceutical compositions of the compound. Ph<sup>1</sup> and Ph<sup>2</sup> are benzene rings with the proviso that carboxy and hydroxy are ortho to one another. Het includes an optionally substituted heterocyclic ring which includes conjugated double bonds and binds to nitrogen in NR. The compound is characterized in that A is a bridge which is stable against reduction because it is not azo, and in that R is hydrogen or lower alkyl. The invention also includes the preparation of the compound and its use as a drug, particularly for treating autoimmune diseases.</p>			



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

Novel Substituted Salicylic Acids

It has long been known that compounds which include a salicylic acid structure coupled by a reductive cleavable bridge -N=N- (=A-) to a ring position in a benzenesulphonic acid which, in turn, is amidated at the amine group of an amine-substituted heterocyclic ring which includes conjugated double bonds can exhibit a therapeutic effect against ulcerative colitis, rheumatoid arthritis and other diseases that are considered to be autoimmune. See for instance Askelöf et al, US-A 2 396 145, Agback and Lindblom, US-A-3 915 951, T. Pullar: Br. J. Clin. Pharmacol., vol. 30, pages 501-510 (1990), I.G. Tumanova, Ia A. Sigidin: Ter-Ark., vol. 59, pages 80-83 (1987), S. Pacheco, K. Hillier, C.L. Smith, Brazilian J. Med. Biol. Res., vol. 23, pages 1323-34 (1990) and C. Astbury, J. Hill, J.R. Lowe, D. Campbell, H.A. Bird, British J. Rheumatology, vol. 29, pages 465-467 (1990). Autoimmune diseases are described and exemplified by, inter alia, I. Roitt, J. Brostoff, D. Male: Immunology, 2nd ed., 1989.

The best known compound possessing the aforesaid therapeutic effect is sulfasalazine. Although analogues of sulfasalazine have been described, none has yet resulted in an accepted drug.

Sulfasalazine is clinically effective in the treatment of autoimmune diseases, such as for instance rheumatoid arthritis, ulcerative colitis, ankylosis spondylitis, reactive arthritis, psoriasis and psoriasis arthritis. See for instance D.R. Porter and H.A. Capell: Bailliers Clin. Rheumatol., vol. 4, pages 535-551 (1990), M.B. Ferraz, P. Tugwell, C.H. Goldsmith and E. Atra: J. Rheumatol., vol. 17, pages 1482-1486 (1990), A.K. Gupta, C.N. Ellis, M.T. Siegel, E.A. Duell, C.E.M. Griffiths, T.A. Hamilton, B.J. Nickoloff and J.J. Voorhees: Arch. Dermatol, vol. 126, pages 487-493 (1990), G. Watkinson: Drugs, vol. 32: suppl. 1, pages 1-11 (1986).

The drugs mostly prescribed in the treatment of rheumatoid arthritis are the antiinflammatory drugs, so-called NSAIDs (Nonsteroidal Anti-Inflammatory Drugs). These drugs are considered to act against the symptoms of the disease. On the other hand, sulfasalazine is classified as a disease-modifying drug. Also belonging to the same group of drugs are gold salt, penicillamine, chloroquine and the immunosuppressive drugs methotrexate, azathioprine and cyclophosphamide, all having radically divergent structures.

A number of pharmacological model systems are available for evaluating drugs which are potentially usable for the treatment of autoimmune diseases. One of the most important properties of the NSAIDs in such model systems is their ability to inhibit prostaglandin synthesis and therewith associated biological effects.

5

Sulfasalazine deviates radically from the active pattern of NSAIDs in such model systems, both clinically and experimentally. Sulfasalazine has been characterized in a number of models judged to be relevant to describe the disease-modifying effective component. For example, it has been shown that the activation of

10 immunocompetent and inflammatory cells are influenced by salphasalazine because it inhibits activation and proliferation of T-lymphocytes and also activation of granulocytes and liberation of mediators.

The clinical effect of sulfasalazine when treating ulcerative colitis has been attributed


15 to 5-aminosalicylic acid, which is formed by reductive cleaving of the mother molecule in the colon. Sulfapyridine is formed at the same time. When treating other autoimmune diseases, such as rheumatoid arthritis for instance, the effective component has been considered to be the intact sulfasalazine molecule or released sulfapyridine or, most probably, both components. The secondary effects of

20 sulfasalazine have essentially been considered to be related to liberated sulfapyridine.

We have now discovered a novel type of analogues to sulfasalazine which exhibit good biological availability and effects in models used in studying autoimmune

25 diseases, such as rheumatoid arthritis and ulcerative colitis for instance. The analogues have the aforescribed structure, but with the azo group (A) replaced with a bridge which is stable against hydrolysis and/or reduction in vivo, for instance in the colon. Such compounds are novel. Thus, in its broadest concept, the invention includes compounds of the structure I

30



and tautomeric forms, salts and solvates, and optionally alkyl esters with 1-6 carbon

35 atoms in the carboxyl group.


In the above structure I, Het is a heterocyclic ring according to the foregoing; Ph<sup>1</sup> is

a benzene ring;  $\text{Ph}^2 (\text{COOH})(\text{OH})$  is a benzene ring with carboxy and hydroxy ortho to one another; R is hydrogen or a lower alkyl ( $\text{C}_{1-6}$ ). Het,  $\text{Ph}^1$ ,  $\text{Ph}^2 (\text{COOH})(\text{OH})$  and the bridge A may be substituted.

- 5 The compounds are characterized in that A is a bridge which is stable against hydrolysis and/or reduction in biological systems. A is preferably a straight carbon chain having at most three carbon atoms ( $-\text{C}-\text{C}-\text{C}-$ ) which includes a carbon-carbon single, double or triple bond, optionally together with an oxo-substituent ( $=\text{O}$ ) on one of the carbon atoms in the chain. By stable against hydrolysis and/or reduction is  
10 meant that A lacks a nitrogen-nitrogen bond, such as azo, and optionally also hydrolysis labile ester and amide bonds as a linking structure between  $\text{Ph}^1$  and  $\text{Ph}^2$ .

- In specific embodiments,  $\text{Ph}^1 = 1,4$  - or  $1,3$  - substituted benzene and  
15  $\text{Ph}^2 (\text{COOH})(\text{OH})$  is an ortho-carboxy-hydroxy substituted phenyl, which may optionally be further substituted with halogen or lower alkyl ( $\text{C}_1-\text{C}_6$ ), preferably methyl, in its 3, 4 or 6 position. -A- is  $-\text{C}\equiv\text{C}-$ ,  $-\text{CH}=\text{CH}-$ ,  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CO}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CHCO}-$ ,  $-\text{CH}_2\text{CO}-$ .

- In specific embodiments of the inventive compounds, the heterocyclic ring in Het is  
20 five-membered or six-membered having two and three conjugated double bonds respectively. The heterocyclic ring in Het may be included in a monocyclic or bicyclic structure. Preferably, Het- is



where Het'- is

25



wherein the free valency in  $\text{N}=\text{C}$ -binds to NR;  
and X is:

30

- (i)  $-\text{N}=\text{CH}-\text{NH}-$ ,  $-\text{N}=\text{CH}-\text{S}-$ ,  $-\text{N}=\text{CH}-\text{O}-$ ,  $-\text{NH}-\text{N}=\text{CH}-$ ,  $-\text{O}-\text{CH}=\text{CH}-$ ,  
 $-\text{CH}=\text{CH}-\text{O}-$ ,  $-\text{NH}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CH}-\text{NH}-$ ,  $-\text{CH}=\text{CH}-\text{S}-$ ,  
 $-\text{CH}=\text{N}-\text{NH}-$ , or

35

- (ii)  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CH}-\text{CH}=\text{N}-$ ,  $-\text{CH}=\text{N}-\text{CH}=\text{CH}-$ ,  
 $-\text{CH}=\text{CH}-\text{N}=\text{CH}-$ ,  $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$ ,

wherein mutually adjacent hydrogen atoms shown in bold (H) can be substituted in pairs with  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ , so as to form a bicyclic structure.

R is hydrogen or lower alkyl ( $\text{C}_{1-6}$ ), preferably hydrogen or methyl.

$\text{R}_1, \text{R}_2, \text{R}_3$  are substituents on carbon atoms in Het'. The groups may be hydrogen, lower alkyl ( $\text{C}_{1-6}$ ), halogen, hydroxy, cyano, carboxy, lower alkoxy ( $\text{C}_{1-6}$ ), benzyl-oxy, lower acyl ( $\text{C}_{1-7}$ ), including acetyl, benzoyl, phenyl, benzyl, etc., wherein any benzene rings that occur may be substituted.

Throughout the descriptive part and the Claims of this specification, by lower alkyl and lower acyl is meant groups which contain 1-6 and 1-7 carbon atoms respectively, optionally with substituents of the aforesaid kind.

A is  $-\text{C}\equiv\text{C}-$ , optionally lower alkyl substituted trans or cis- $\text{CH}=\text{CH}-$ ,  $-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CO}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CH}-\text{CO}-$ ,  $-\text{CO}-$ ,  $-\text{CH}_2-\text{CO}-$ ,  $-\text{CH}_2-$ , preferably  $-\text{C}\equiv\text{C}-$  or trans- $\text{CH}=\text{CH}-$ ;

$\text{Ph}^2$  is  $\text{C}_6\text{H}_2\text{R}_4$ , where  $\text{R}_4$  is hydrogen, halogen or lower alkyl, preferably hydrogen or methyl; and tautomeric forms thereof and salts with alkali metals, preferably sodium, with calcium or magnesium, or with pharmaceutically acceptable amines, such as crystal solvates which include pharmaceutically acceptable solvents, such as water, acetone and ethanol for instance, and also pharmaceutical compositions thereof.

The invention also relates to the use of the compound as a drug, primarily for treating autoimmune diseases in analogy with the use of sulfasalazine and potentially also other inflammatory conditions. Other aspects of the invention include the preparation of the compounds and the production of pharmaceutical compositions containing said compounds and intended for the medicinal indications mentioned below.

The inventive compounds have an immunomodulating effect in biological systems, for instance by inhibiting immunocompetent and inflammatory cell activation and in their pharmacological profile are similar to sulfasalazine but are often more active.


Consequently, the inventive compounds are potential drugs for treating such autoimmune diseases as rheumatoid arthritis, ulcerative colitis, ankylosing spondylitis, reactive arthritis, psoriasis, psoriasis arthritis, Morbus Crohn, multiple

sclerosis, type 1 diabetes, scleroderma, myasthenia gravis, Sjögrens syndrome, systematic lupus erythematosus and chronic asthma. The inventive compounds can also be used conceivably for treating other diseases, particularly those diseases which have an immunological component, irrespective of whether the compounds are known for this medicinal science or not.

Because the novel compounds are unable to form sulfapyridine or any other toxic analog thereof, due to the particular nature of said compounds, it is extremely probable that said compounds lack the side effects due to free sulfapyridine. Our experiments have shown that the biological availability of laboratory animals is often much greater for the inventive compounds than for sulfasalazine and its analogues.

In summary, the novel compounds should prove highly beneficial as drugs for the treatment of autoimmune diseases. The novel inventive compounds can be produced in several ways.

One method involves reacting compound II with compound III.



where R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, Ph<sup>1</sup> and Het' are the same as above, R<sub>5</sub> is hydrogen or preferably lower alkyl having at most six carbon atoms, and X and Y pairwise are -C≡CH and Z<sub>1</sub>-; -Z<sub>1</sub> and HC≡C-; -CH=CH<sub>2</sub> and Z<sub>1</sub>-; -Z<sub>1</sub> and -CH=CH<sub>2</sub>; -CO-CH<sub>3</sub> and HCO-; -COH and CH<sub>3</sub>-CO-; -CH<sub>2</sub>COZ<sub>2</sub> and H-; -CHO and Z<sub>3</sub>CH<sub>2</sub>-; -CH<sub>2</sub>Z<sub>3</sub> and HCO-; wherein Z<sub>1</sub> is bromo or iodo, Z<sub>2</sub> is halo, preferably chloro, and Z<sub>3</sub> is the residue of a phosphororganic compound of the Wittig type, for instance a triphenyl phosphonium group or a dialkyl phosphonate group.

The compounds II and III are coupled together in a known manner, whereafter when R<sub>5</sub> is a lower alkyl, the coupling product is hydrolyzed, preferably in the presence of alkali metal hydroxide, whereafter the product is acidified and then isolated.

When X and Y are -C≡CH and Z<sub>1</sub> or Z<sub>1</sub> and HC≡C-, the aforesaid coupling process is preferably effected by means of the so-called Heck's reaction, implying catalysis of a palladium compound and a copper compound in the presence of a base and a

solvent. Suitable palladium compounds are dichlorobis(triphenylphosphine)-palladium, dichlorobis[tris(2-methylphenyl)]palladium or tetrakis(triphenylphosphine)palladium. A suitable palladium compound may optionally be prepared in situ, by using, e.g., palladium chloride or palladium acetate together with  
5 triphenylphosphine, tris(2-methylphenyl) phosphine or 1,3-bis(diphenylphosphino) propane, etc. Suitable copper compounds are copper (I) iodide or copper (I) bromide. Suitable bases are amines, preferably tertiary amines, such as, for instance, triethylamine or tributylamine or other inorganic or organic bases such as, for instance, sodium hydrocarbonate or sodium acetate. Suitable solvents are, for instance, N,N-dimethylacetamide, N,N-dimethylformamide, ethanol, acetone, tetrahydrofuran,  
10 dioxane, toluene, etc., preferably N,N-dimethylacetamide.

When X and Y are  $-\text{CH}=\text{CH}_2$  and  $\text{Z}_1$  or  $\text{Z}_1$  and  $-\text{CH}=\text{CH}_2$ , the aforesaid coupling is preferably effected by means of the so-called Heck's reaction, implying catalysis of a  
15 palladium compound in the presence of a base and solvent. Suitable palladium compounds are divalent inorganic or organic salts of palladium, e.g. palladium (II) chloride and palladium (II) acetate. The salts may optionally be stabilized with customary ligands such as, for instance, triphenylphosphine in the form, for instance, of dichlorobis(triphenylphosphine) palladium. Alternatively, a 0-valence palladium  
20 compound can be used such as, for instance, bis(dibenzylidene acetone)palladium or tetrakis(triphenylphosphine)palladium. Suitable bases are tertiary amines such as, for instance, triethylamine or inorganic bases such as sodium hydrogen carbonate or sodium acetate. Suitable solvents are, for instance, N,N-dimethylacetamide, N,N-dimethylformamide, toluene, tetrahydrofuran, dioxane,  
25 acetone, etc.

Although the compound of formula  $(\text{R}_1, \text{R}_2, \text{R}_3)\text{-Het}'\text{-NR-SO}_2\text{-Ph}^1\text{-X}$ , where X is ethenyl, can be produced in several ways, each known per se, the compound is preferably produced by reacting a compound of the formula  $(\text{R}_1, \text{R}_2, \text{R}_3)\text{-Het}'\text{-NR-SO}_2\text{-Ph}^1\text{-X}$ , where X is bromo or iodo, with ethylene in the presence of a catalytic  
30 quantity of a palladium compound, a base and a solvent, where catalysts and solvents are of a similar type to those used in the reaction between II and III. Another method involves partially reducing a corresponding ethynyl compound with gaseous hydrogen in the presence of a catalyst appropriate for this purpose.

35

The compound of the formula II, where X is bromine or iodine, is produced in a known manner, for instance by reacting corresponding halogen benzene sulfonyl




chloride with corresponding heterocyclic amine in the presence of a base, for instance pyridine or the heterocyclic amine in excess.

When X and Y in formula II and III respective are -CO-CH<sub>3</sub> and HCO- or  
 5 -CHO and CH<sub>3</sub>-CO-, the aforesaid coupling is effected in a known manner, in the presence of a solvent and a basic or acid condensation agent. Suitable bases are, for instance, sodium hydroxide or potassium hydroxide. Suitable acids are strong mineral acids, such as hydrochloric acid or sulphuric acid. Suitable solvents are water, lower alcohols or mixtures thereof. The compound of formula II, where -X is  
 10 -CO-CH<sub>3</sub> or -CHO is prepared in a known manner, for instance by reacting corresponding formylbenzene sulfonyl chloride or acetylbenzene sulfonyl chloride with corresponding heterocyclic amine, in the presence of a base, for instance pyridine or the heterocyclic amine in excess.

15 When -X and Y- in compound II and III respectively are -CH<sub>2</sub>COZ<sub>2</sub> and H-, the reaction is effected in a known manner under Fridel-Craft's conditions, in the presence of a Lewis acid, preferably aluminium chloride in an inert solvent, such as chlorinated hydrocarbon for instance. The compound II, where -X is -CH<sub>2</sub>COZ<sub>2</sub>, is prepared in a known manner, for instance by reacting corresponding alkoxycarbonyl-  
 20 benzenesulfonyl chloride with corresponding heterocyclic amine in the presence of a base, for instance pyridine or the heterocyclic amine in excess, whereafter the intermediary amine is hydrolyzed to corresponding carbonic acid and the acid is halogenated in a known manner, for instance by reacting with thionylchloride or a phosphorous halogenide.

25

Another method is by reacting a compound of formula IV with a compound of formula V




30

where Het', Ph<sup>1</sup>, R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> have the same significance as above and R<sub>6</sub> is a lower acyl, preferably acetyl or the residue of an aliphatic or aromatic sulfonic acid, preferably methyl sulfonyl or p-toluenesulfonyl; in the presence of  
 35 a basic condensation agent and a solvent, wherein the basic condensation agent may be the compound IV in excess or preferably an organic base, for instance pyridine, whereafter the resultant intermediary compound is subjected to base-catalyzed

hydrolysis, preferably by reaction with alkali metal hydroxide in water.

The aforesaid compound of formula V is produced in a known manner, by reacting corresponding sulfonic acid or its alkali metal sulfonate with a suitable chlorinating agent, preferably thionylchloride in the presence of N,N-dimethylformamide. The sulfonic acid is produced in a known manner according to any one of the abovementioned general methods, preferably by converting corresponding ethynyl or ethenyl aryl compound with an aryl iodide or aryl bromide. A third method of producing a compound according to the invention is by transforming the bridge A in an inventive compound or its intermediate having the formula VI (below) to another bridge in a known manner, optionally with subsequent transformation of the intermediate to the final compound.

One example of such methods involves the addition of water to a compound of formula VI



where R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, Het', Ph<sup>1</sup>, R<sub>4</sub> are the same as above, R<sub>7</sub> is hydrogen or R<sub>5</sub>, where R<sub>5</sub> is the same as above, R<sub>8</sub> is hydrogen or R<sub>6</sub>, where R<sub>6</sub> has the same meaning as above, and -A- is -C≡C-. The resultant compound has the formula VI with A=-CH<sub>2</sub>CO-. Addition of water to the triple bond takes place in the presence of a mineral acid or a strong organic acid, preferably formic acid, optionally in the presence of metal salts of, e.g., mercury or palladium, with subsequent alkaline hydrolysis when R<sub>7</sub> and/or R<sub>8</sub> are not hydrogen. Another method is to reduce the compound of formula VI, where R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, Het', Ph<sup>1</sup>, R<sub>4</sub>, R<sub>7</sub> and R<sub>8</sub> are the same as above and -A- is -CH<sub>2</sub>CO- to a corresponding compound in which -A- is -CH<sub>2</sub>CHOH-, and by then converting this compound with mineral acid so as to eliminate water and form a compound where -A- is -CH=CH- with subsequent alkaline hydrolysis when R<sub>7</sub> and/or R<sub>8</sub> are not hydrogen.

A third method is to reduce a compound of formula VI, where R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, Het', Ph<sup>1</sup>, R<sub>4</sub>, R<sub>7</sub> and R<sub>8</sub> are the same as above and -A- is -C≡C- or -CH=CH- to a corresponding compound in which -A- is -CH<sub>2</sub>CH<sub>2</sub>- by means of catalytic hydrogenation in a known manner, followed by alkaline hydrolysis when R<sub>7</sub> and/or R<sub>8</sub> are not hydrogen. When the inventive compound is a salt, the compound is

produced, for instance, by first producing a corresponding acid and then reacting this acid with a corresponding salt former, for instance in the form of a metal hydroxide or organic amine, in the presence of a solvent, preferably water or a mixture of water and one or more organic solvents, or optionally solely in the presence of an organic solvent. When the salt is not readily soluble in the solvent used, the salt is preferably allowed to crystallize from the solvent and is isolated by filtration or some similar process. When the not readily dissolved salt is an alkali metal salt, preferably sodium or potassium, the final hydrolysis applied to remove any protective groups may be carried out so that the synthesis mixture will be neutralized and the desired salt is crystallized directly from the mixture. If the salt is relatively soluble in the solvent used, the salt is preferably produced by reacting equimolar quantities, whereafter the solvent is removed by evaporation.

It is possible in some instances for a compound to form a distinct solvate with a solvent. Such solvates, which contain pronounced quantities of solvent, are also included by the invention when the solvents used are pharmaceutically acceptable, such as water, acetone and ethanol for instance. Such a solvate is produced by crystallization from the solvent concerned, optionally in mixture with other solvents. Optionally, crystallization can be effected by acidifying a soluble salt of the compound in the presence of the solvate-forming solvent.

The invention also embraces pharmaceutical compositions which include the compound of formula I, intended particularly for oral use, optionally in combination with an organic or inorganic inert carrier suitable for oral ingestion and/or other conventional additives. The pharmaceutical composition may, for instance, be in tablet, dragee, capsule form, etc., optionally enteric coated, or solutions and suspensions containing inventive compounds. The pharmaceutical composition can be produced in a known manner by a person competent within this field, by mixing the compound or formula I with the desired carrier material and/or further additives, and converting this mixture to a desired galenic form in accordance with the foregoing. Solutions and suspensions are prepared in a known manner with the aid of pharmaceutically appropriate additives. The dosage is adapted to requirements and wishes in individual situations, although dosages of 50-2000 mg/day in the case of adult patients can be mentioned as a general indication.

## Examples.

- 5 The identity of all final compounds have been assured by their NMR spectra and the purity by thin layer chromatography (TLC) or liquid chromatography. NMR proton spectra have been determined on a 500 MHz NMR spectrometer, with deuterated dimethylsulfoxide as solvent unless otherwise indicated. NMR chemical shift are given in ppm.

10

## Example 1

2-Hydroxy-5-[[4-[(2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoic acid.

## 15 Example 1a

*Methyl 5-[(trimethylsilyl)ethynyl]-2-hydroxybenzoate*

- To a solution of methyl 2-hydroxy-5-iodobenzoate (275 g, 0.92 mol) in dry triethylamine (2000 ml) was added dichlorobis(triphenylphosphine)palladium (3 g, 0.004 mol) and copper(I) iodide (1.6 g, 0.008 mol). The mixture was deaerated with nitrogen. Trimethylsilylacetylene (100 g, 1.0 mol) was added with a syringe and the reaction mixture was heated to 50°C. After 30 minutes a voluminous precipitate of amine hydroiodide had formed and after 2 h the reaction was complete according to TLC. The mixture was filtered and the filtrate was evaporated *in vacuo*. The residue was purified by dry-flash chromatography on silica, using toluene as the eluent. Yield 211 g (92 %).

25

## Example 1b

*Methyl 5-ethynyl-2-hydroxybenzoate.*

- 30 A mixture of methyl 5-[(trimethylsilyl)ethynyl]-2-hydroxybenzoate (100 g, 0.57 mol) and potassium fluoride dihydrate (150 g, 1.59 mol) in DMF (600 ml) was stirred for 4 h at room temperature. The solution was extracted with ether (3 x 400 ml) and the combined ether extracts were washed with 1 M HCl (2 x 200 ml) and water (2 x 100 ml). The ether layer was dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. Yield 66.5 g (93 %).

35

## Example 1c

*4-Iodo-N-(2-pyridinyl)benzenesulfonamide.*

4-Iodo-benzenesulfonyl chloride, (52.3 g, 0.17 mol) was dissolved in dichloromethane (300 ml) and 2-aminopyridine (65 g, 0.69 mol) was added. The solution was stirred at room temperature for 3 days, washed with 2 M sulfuric acid (2 x 200 ml) and water (100 ml), dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. Yield 46.9 g (76 %).

## Example 1d

*Methyl 2-hydroxy-5-[[4-[(2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoate*

4-Iodo-N-(2-pyridinyl)benzenesulfonamide (35 g, 97 mmol) was dissolved in a mixture of triethylamine and tetrahydrofuran (750 + 750 ml). The mixture was deaerated with nitrogen and dichlorobis(triphenylphosphine)palladium (1.2 g, 1.7 mmol) and copper(I) iodide (0.6 g, 3.4 mmol) were added. Finally methyl 5-ethynyl-2-hydroxybenzoate (23 g, 130 mmol) was added. The mixture was heated to 60°C for 4 h, evaporated to dryness. The residue was dissolved in chloroform (1000 ml) and washed with water (3 x 200 ml) and dried with MgSO<sub>4</sub>. The solvent was evaporated *in vacuo*. Yield 31 g (77 %). The product can be purified by flash-chromatography on silica with chloroform as the eluent.

## Example 1e

*2-Hydroxy-5-[[4-[(2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoic acid*

The ester from Example 1d (23.7 g, 58 mmol) was dissolved in 1 M NaOH (190 ml) and refluxed for 6 h. The cooled solution was acidified with an excess of 6 M HCl. The precipitated formed was allowed to stand for 2 h, filtered off and washed with water (3 x 200 ml). The white product was dried at 60°C *in vacuo*. Yield 21.7 g (95 %).

<sup>1</sup>H-NMR: Spin system A: δ 7.00(d), 7.66(dd), 7.95(d): (=salicylate ring).  
Spin system B: δ 7.66(d,2H), 7.87(d,2H): (=central benzene ring)  
Spin system C: δ 7.18(d), 7.73(ddd), 6.84(dd), 7.96(dd): (=pyridine ring)

## Example 2.

2-Hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)-sulfonyl]phenyl]ethynyl]benzoic acid.

5

## Example 2a

*4-Iodo-N-(3-methyl-2-pyridinyl)benzenesulfonamide.*

10

This compound was prepared essentially as the corresponding compound in Example 1c. Yield 76 %.

## Example 2b

*Methyl 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoate.*

15

This compound was prepared essentially as the corresponding compound in Example 1d. Yield 76 %

## Example 2c

20 *2-Hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoic acid.*

This compound was prepared essentially as the corresponding compound in Example 1e. Yield 24.7 g (85 %).

25

<sup>1</sup>H-NMR: Spin system A: δ 7.04(d), 7.70(dd), 8.00(d): (=salicylate ring).  
Spin system B: δ 7.70(d,2H), 7.96(d,2H): (=central benzene ring)  
Spin system C: δ 7.66(d, broad), 6.83(s, broad), 7.84(s, broad):  
(=pyridine ring).  
Spin system D: δ 2.15(s): (=methyl group)

30

The broadening of signals in the pyridine ring is typical for 3-methyl substituted derivatives.

## Example 3.

2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.

## 5 Example 3a

*Methyl 5-ethenyl-2-hydroxybenzoate.*

Methyl 5-ethynyl-2-hydroxybenzoate (8 g, 45.4 mmol), prepared according to Example 1b, was dissolved in a mixture of diethyl ether and light petroleum (200 +  
10 200 ml) in a hydrogenation flask. Quinoline (1.7 ml) and palladium on CaCO<sub>3</sub>, poisoned with Pb (Lindlar catalyst; 200 mg) were added, and the flask was attached to an atmospheric pressure hydrogenation apparatus. The mixture was stirred for 2 h at room temperature, by then the calculated amount of hydrogen had been consumed. The catalyst was filtered off and the solution was evaporated to dryness. The product  
15 was purified by flash column chromatography on silica, using toluene as an eluent. A nearly quantitative yield was obtained.

## Example 3b

*Methyl 2-hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoate.*

20

4-Iodo-N-2-pyridinylbenzenesulfonamide (25 g, 69 mmol) prepared according to Example 1c, and methyl 5-ethenyl-2-hydroxybenzoate (13.5 g, 76 mmol) were dissolved in a mixture of triethylamine and tetrahydrofuran (150 + 800 ml). The mixture was heated to 80° for 18 h. Palladium(II) acetate (in a total of 0.5 g) was  
25 added in small portions during the whole reaction time. The amine hydroiodide formed was filtered off and the filtrate was evaporated to dryness. The residue was dissolved in chloroform (500 ml) and purified by flash-chromatography on silica with chloroform as the eluent. The solvent was evaporated *in vacuo* and the residue was dissolved in the smallest possible amount of hot tetrahydrofuran. The product  
30 was precipitated by addition of diethyl eter. Yield 13.5 g (47 %).

## Example 3c

*2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.*

35 Methyl 2-hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoate (12 g, 29 mmol) was dissolved in 1 M KOH (120 ml) and the solution was refluxed for 7 h. The cooled solution was acidified with an excess of 1 M hydrochloric acid. The

precipitate was filtered off and washed with water (3 x 200 ml). The product was redissolved in a small portion of 1 M NaOH. Water and dioxane (500 + 500 ml) was added and the solution was heated to 100°C. The solution was acidified with an excess of 1 M HCl, and the product precipitated upon cooling. Yield 9 g (75 %).

5

<sup>1</sup>H-NMR: Spin system A: δ 7.02(d), 7.84(dd), 8.04(d): (=salicylate ring).  
Spin system B: δ 7.73(d,2H), 7.88(d,2H): (=central benzene ring)  
Spin system C: δ 7.22(d), 7.74(m), 6.89(ddd), 8.04(m): (=pyridine ring)  
Spin system D: δ 7.19(d), 7.38(d): (=ethenediyl bridge)

10

#### Example 4.

2-Hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.

15

#### Example 4a

*Methyl 2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)-sulfonyl]phenyl]ethenyl]benzoate.*

20

This compound was prepared analogously to Example 3 from 4-iodo-N-(3-methyl-2-pyridinyl)-benzenesulfonamide, prepared according to Example 2a, and methyl 5-ethenyl-2-hydroxybenzoate, prepared according to Example 3a. Yield 52 %.

#### Example 4b

25

2-Hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.

This compound was prepared analogously to Example 3. Yield 82 %.

#### <sup>1</sup>H-NMR.

30

Spin system A: δ 7.02(d), 7.85(dd), 8.04(d): (=salicylate ring).  
Spin system B: δ 7.74(d,2H), 7.92(d,2H): (=central benzene ring)  
Spin system C: δ 7.63(d, broad), 6.81(broad), 7.85(broad): (=pyridine ring)  
Spin system D: δ 7.21(d), 7.39(d): (=ethenediyl bridge)  
Spin system E: δ 2.15(s): (=methyl group)

35

The broadening of signals in the pyridine ring is typical for 3-methyl substituted derivatives.



## Example 5

Isobutyl 2-hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoate.

## 5 Example 5a

*Isobutyl 2-hydroxy-5-iodobenzoate.*

2-hydroxy-5-iodobenzoic acid (477 g, 1.8 mol), isobutanol (225 g), conc sulfuric acid (20 ml) in toluene (2 l) was refluxed for 6 h using a water separator. The solution was cooled and washed with water. The solution was treated with decolorizing carbon and the toluene was evaporated. The residue was recrystallized from methanol. Yield 376 g.

## Example 5b

15 *Isobutyl 5-ethenyl-2-hydroxybenzoate.*

Isobutyl 2-hydroxy-5-iodobenzoate (16 g, 0.05 mol), tributylamine (13.9 g, 0.075 mol), and palladium(II) acetate (0.11 g, 0.5 mmol) were dissolved in dimethylacetamide (100 ml) and the solution was charged into a 250 ml pressure-reactor.

Air was expelled by inlet of argon at 10 bar followed by venting to ambient pressure. This procedure was repeated twice. The temperature was increased to 90 °C. Ethene at 13 bar was let in. After stirring for 5 min the gas was vented and the ethene-pressure was again raised to 13 bar. The temperature was increased to 110 °C during 30 min with vigorous stirring and continuous inlet of ethene to keep the pressure constant at 13 bar. After 1.5 h the temperature was decreased to 20 °C and the ethene was removed by means of argon inlet.

30 The solvent was evaporated at 70 °C. Water (150 ml) was added. The pH was adjusted to about 5 with hydrochloric acid and the solution was extracted with toluene (2x50 ml). After reextraction with dilute sodium hydrogencarbonate (50 ml) the combined toluene extracts were evaporated to leave the product as an oil. Distillation at 0.5 mm Hg and 100-105 °C afforded 15.4 g (70 %) of chromatographically pure material. During distillation the material was protected from polymerisation by addition of 10 mg of hydroquinone.

## Example 5c

*Isobutyl 2-hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoate.*

Palladium (II) acetate (0.15 g, 0.67 mmol) was added to a stirred solution of isobutyl 5-ethenyl-2-hydroxybenzoate (6.6g, 30 mmol) and 4-iodo-N-2-pyridinylbenzenesulfonamide (10.8 g, 30 mmol) in dimethylacetamide (40ml) and tributylamine (10 ml, 42 mmol) at 95 °C. After 100 min. the solution was filtered and evaporated at 50-60 °C (bath temp.) until solid material appeared. The mixture was diluted with isopropanol (0.5 l) and chilled in a freezer. The product was filtered off to yield 7.1g (52 %).

<sup>1</sup>H-NMR.

Spin system A: δ 7.0(d), 7.84(dd), 7.93(d): (=salicylate ring).

Spin system B: δ 7.70(d,2H), 7.84(d,2H): (=central benzene ring)

Spin system C: δ 7.18(d), 7.69(ddd), 6.85(ddd), 8.0(dd): (=pyridine ring)

Spin system D: δ 7.12(d), 7.36(d): (=ethenediyl bridge)

Spin system E: δ 1.0(d, 6H), 2.1(m, 1H), 4.1(d,2H): (=isobutyl group).

## Example 6

2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.

## Example 6a

*Isopropyl 2-hydroxy-5-iodobenzoate*

2-Hydroxy-5-iodobenzoic acid (184 g, 0.7 mol), isopropanol (50 ml), conc sulfuric acid ( 5 ml) and p-toluenesulfonic acid (10 g) in toluene (1 l) was refluxed using a water separator. Portions of isopropanol was added intermittently to compensate for loss of 2-propanol by dehydration. After 24 h, the solution was cooled and washed with water. The toluene was evaporated partially. Iso-octane was added, and precipitated 2-hydroxy-5-iodobenzoic acid (106 g) filtered off. The solution was taken to dryness and the residue recrystallized from methanol. Yield 64 g, (71 %, corrected for recovered starting material).

## Example 6b

*Isopropyl 5-ethenyl-2-hydroxybenzoate.*

5 Isopropyl 2-hydroxy-5-iodobenzoate (30.6 g, 0.1 mol) was dissolved in a mixture of tributylamine (27.8 g, 0.15 mol) and dimethylacetamide. After addition of palladium(II) acetate and air venting the temperature was increased to 110 °C under argon. The ethene inlet was carried out following the procedure of Example 5b.

10 The reaction was arrested after 1.5 h by cooling to room temperature and simultaneous venting of the ethene.

After evaporation of the solvents in vacuo at 70 °C water (100 ml) was added and pH was adjusted to about 5 with hydrochloric acid. Extraction with 2x50 ml toluene and subsequent evaporation afforded an oily raw material.

15 The product was distilled at 0.7 mm Hg and 95- 98 °C. Yield 15.5 g (75 %) of chromatographically pure material.

## Example 6c

20 *2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.*

Isopropyl 5-ethenyl-2-hydroxybenzoate was reacted with 4-iodo-N-(2-pyridinyl)benzenesulfonamide similarly to the procedure of Example 5c. After the reaction, the solvent was evaporated, toluene was added and the suspension filtered and washed with toluene. After drying on the filter and washing with water the material was hydrolyzed and precipitated similarly to the procedure of Example 3c. Yield 40 %

NMR proved the identity with the product of Example 3.

## Example 7

2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.

- 5 Isobutyl 2-hydroxy-5-iodo-benzoate (16 g, 0.05 mol) was reacted with ethene as described in Example 5b

The reaction mixture was cooled to 85 °C and 4-iodo-N-(2-pyridinyl)benzenesulfonamide (14.8 g, 0.0425 mol.) and palladium(II) acetate (0.11 g, 0.5 mmol) were added with stirring.

10

After 4 hours at 85 °C, all the sulfonamide had been consumed but the reaction mixture still contained some isobutyl 5-ethenyl-2-hydroxybenzoate. A further amount of sulfonamide (2.6 g, 0.0075 mol) was added and the reaction was continued for 18 h.

15

After addition of acetic acid the hot mixture was poured into boiling isopropanol (450 ml). Upon cooling slowly to 4 °C the solution yielded 11.3 g of the product. A further 1.5 g was obtained by concentrating the mother liquor to about 100 ml and cooling to 4° C. Total yield 12.7 g (56 %)

20

The material was hydrolyzed overnight in refluxing 1M potassium hydroxide (150 ml) and the product was precipitated by addition of an excess of 1M HCl. Recrystallisation from formic acid/water (70/30) yielded after filtration and drying 8.5 g (43 %) product.

25

NMR proved the identity with the product of Example 3.

## Example 8

30 Isobutyl 2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]-ethenyl]benzoate.

10 % Palladium on charcoal (0.5g) was added to a solution of isobutyl 5-ethenyl-2-hydroxybenzoate (6.6g, 30mmol) and 4-iodo-N-(3-methyl-2-pyridinyl)benzenesulfonamide (11.2 g, 30 mmol) in dimethylacetamide (10 ml) and tributylamine (10 ml, 42 mmol) at 110 °C. After 90 min. the solution was filtered and diluted with isopropanol (0.3 l) and chilled in a freezer. The product was filtered

35

off, dried and dissolved in boiling formic acid (100 ml). The solution was filtered and water (65 ml) was added. After cooling to refrigerator temperature the product was filtered off and dried to yield 7.2g (51 %) product.

5  $^1\text{H}$ -NMR.

Spin system A:  $\delta$  7.0(d), 7.84(d, broad), 7.94(s, broad): (=salicylate ring).

Spin system B:  $\delta$  7.69(d,2H), 7.9(d,2H): (=central benzene ring)

10 Spin system C:  $\delta$  7.58(broad), 6.74(broad), 7.8-8.0(broad): (=pyridine ring)

Spin system D:  $\delta$  7.12(d), 7.35(d): (=ethenediyl bridge)

Spin system E:  $\delta$  2.1(s, broad): (=methyl group)

Spin system F:  $\delta$  1.0(d, 6H), 2.1(m, obscured by methyl of spin system E), 4.1(d,2H): (=isobutyl group).

15 The broadening of signals in the pyridine ring is typical for 3-methyl substituted derivatives.

Example 9

20 Ethyl 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoate.

Example 9a

*Ethyl 5-ethynyl-2-hydroxybenzoate.*

This compound was synthesized from ethyl 2-hydroxy-5-iodobenzoate analogously to the method 1a and 1b for the corresponding ethyl ester.

25

Example 9b

*Ethyl 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoate.*

30 Tetrakis(triphenylphosphine)palladium (0.29 g, 0.25 mmol) and copper(I)iodide (0.095 g, 0.5 mmol) were added to a solution of ethyl 5-ethynyl-2-hydroxybenzoate (10.5g, 55 mmol) and 4-iodo-N-(3-methyl-2-pyridinyl)benzenesulfonamide (18.7 g, 50 mmol) in tetrahydrofuran (75 ml) and triethylamine (75 ml) at 55 °C. After 2 h the solid product was collected with filtration and washed with hydrochloric acid (1M). The material was boiled in formic acid (300 ml) for 15 min. and filtered.  
35 Water (200 ml) was added and the product crystallized upon cooling. Yield 17.3 g (79 %).

<sup>1</sup>H-NMR (Pyridine-d<sub>5</sub> as solvent)

Spin system A: δ 7.11(d), 7.70(dd), 8.14(d): (=salicylate ring).

Spin system B: δ 7.71(d,2H), 8.33(d,2H): (=central benzene ring)

5 Spin system C: δ 7.37(d), 6.64(dd), 7.94(d): (=pyridine ring)

Spin system D: δ 2.11(s, broad): (=methyl group)

Spin system E: δ 1.22(t, 3H), 4.33(q, 2H): (= ethyl group).

#### Example 10

10 2-Hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]-benzoic acid.

#### Example 10a

15 4-(3-Hydroxy-3-methyl-1-butyne-1-yl)-N-(3-methyl-2-pyridinyl)benzenesulfonamide.  
4-Iodo-N-(3-methyl-2-pyridinyl)benzenesulfonamide (37.4 g, 0.1 mol) was added in portions to a mixture of triethylamine (31 ml, 0.22 mol) and dimethylacetamide (25 ml) at 60 °C. Dichlorobis(triphenylphosphine)palladium (0.14 g, 0.002 mol) and CuI (0.08 g, 0.004 mol) were added under a nitrogen blanket. Finally, 2-methyl-3-butyne-2-ol (10.4 g, 0.125 mol) was added in portions during 20 minutes in order to keep the  
20 temperature of the exothermal reaction at 65-70 °C.

The mixture was stirred at 65 °C for 1 hour. Water (250 ml) was added and after cooling to room temperature a cloudy precipitate was filtered off. 85 % potassium hydroxide (13.2 g 0.2 mol) dissolved in water (100 ml) was added. About 100 ml of  
25 water/triethylamine was distilled off in vacuo at 50 °C. The remaining solution was extracted with isobutylmethylketone (3x30 ml) and from the water-phase about 50 ml of isobutylmethylketone/water was evaporated at 50 °C.

The volume was adjusted to 300 ml with water and ethanol (100 ml ) was added.  
30 Hydrochloric acid was cautiously added dropwise at 60 °C until the pH reached 4. Upon cooling to 8 °C the product precipitated. The product was filtered off, washed with water and dried. Yield 29.6 g, (90 %).

#### Example 10b

35 4-Ethynyl-N-(3-methyl-2-pyridinyl)benzenesulfonamide.  
4-(3-Hydroxy-3-methyl-1-butyne-1-yl)-N-(3-methyl-2-pyridinyl)benzenesulfonamide (49.6 g 0.15 mol) and water (900 ml) was heated to about 90 °C under a nitrogen

blanket. 85 % potassium hydroxide (64.9 g, 1.05 mol) was added at a rate sustaining vigorous boiling. A vigorous boiling was maintained for 2 hours without reflux in order to efficiently remove the acetone formed in the reaction. Water was added to keep the volume approximately constant.

5

The hot solution was added to ethanol (200 ml) containing a small amount of SO<sub>2</sub>. Hydrochloric acid was added dropwise at 90 °C. When an opalescence was observed, activated carbon (1.5 g) was added and the solution filtered. Continued addition of hydrochloric acid was maintained to pH 4 causing precipitation of the product.

10

After cooling to 8 °C the solid material was filtered off and washed with water. The product was dried at 80 °C in vacuo for 18 h. Yield 38.2 g (94 %).

#### Example 10c

15 *Ethyl 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoate.*

Dichlorobis(triphenylphosphine)palladium (0.07 g, 0.1 mmol) and copper(I)iodide (0.04 g, 0.2 mmol) were added to a solution of ethyl 2-hydroxy-5-iodobenzoate (14.9 g, 50 mmol) and 4-ethynyl-N-(3-methyl-2-pyridinyl)benzenesulfonamide (14.3 g, 50 mmol) in triethylamine (10 g, 0.1 mol) and dimethylacetamide (60ml) at 65 °C. Hot ethanol was added after 1h and then hot water (100 ml) with stirring. The solid was filtered off after cooling in the refrigerator and recrystallized from formic acid and water to yield 17.0 g (77 %) product.

#### 25 Example 10d

*2-Hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoic acid*  
Ethyl 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoate (15.5g, 35 mmol) was refluxed in potassium hydroxide (10 g, 0.15 mol) in water (100 ml) for 1 h. After cooling to ca 70 °C, ethyl acetate (50 ml) was added and hydrochloric acid was added to reach pH 7-8. Cooling during stirring and then filtration gave a solid that was dissolved in acetone (100 ml) and water (100 ml). The solution was acidified at ca 50 °C and cooled, filtered and washed with water. Yield 8.0 g (56 %).

35 NMR proved the identity with the product of Example 2.

## Example 11

Potassium 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]-ethynyl]benzoate.

## Example 11a

5     4-(3-Hydroxy-3-methyl-1-butyn-1-yl)-N-(3-methyl-2-pyridinyl)-benzenesulfonamide.

The synthesis was performed as in Example 10a in 0.2 mole scale with acetone (100 ml) instead of dimethylacetamide as solvent. The reaction time at reflux (60-62 °C) was 2 h.

10     Yield 60.0 g, 91 %.

## Example 11b

4-Ethynyl-N-(3-methyl-2-pyridinyl)benzenesulfonamide.

This synthesis was run exactly as in Example 10b.

15

## Example 11c

Ethyl 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]-benzoate

20     The synthesis was carried out as in Example 10c, except that hydrochloric acid was added after the ethanol and water addition. Yield 72 %.

## Example 11d

Potassium 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]-phenyl]ethynyl]benzoate

25     Ethyl 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoate (15.5g, 35 mmol) was refluxed in potassium hydroxide (10 g, 0.15 mol) in water (100 ml) for 1 h. After cooling to ca 70 °C, ethyl acetate (50 ml) was added and hydrochloric acid was added to reach pH 7-8. Cooling with stirring and then filtration gave directly 8.5 g of the product (51  
30     %).

The corresponding acid of the product proved to be identical with the product of Example 2.



## Example 12.

2-Hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.

5

## Example 12a

*4-(2-Bromoethyl)-N-(3-methyl-2-pyridinyl)benzenesulfonamide*

3-Methyl-2-pyridinamine (431 g, 4.0 mol) was dissolved in dichloromethane (700 ml) and 4-(2-bromoethyl)benzenesulfonyl chloride (340 g, 1.33 mol) was added at room temperature. After 1 h at 20 °C, the solution was heated to 35 °C for 2 h. Excess dilute hydrochloric acid was added and the phases separated. After washing once with water, the organic phase was dried with MgSO<sub>4</sub> and treated with decolorizing carbon. The solvent was evaporated and warm methanol added to the residue, which crystallized. The material was collected and recrystallized from toluene. Yield 178 g. A further crop of 36 g was obtained by workup of the mother liquor. Total yield 45 %.

10  
15

## Example 12b

*4-Ethenyl-N-(3-methyl-2-pyridinyl)benzenesulfonamide.*

20

4-(2-Bromoethyl)-N-(3-methyl-2-pyridinyl)benzenesulfonamide (183 g, 0.5 mol) and potassium hydroxide (120 g, 1.83 mol) and hydroquinone (1 g) in ethanol (2 l) was refluxed for 1 h. 5 l water was added and the solution acidified with acetic acid. The precipitated product was collected by filtration, washed with water and dried. Yield 137 g (quantitative).

25

## Example 12c

*Isobutyl 2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]-ethenyl]benzoate.*

30

Isobutyl 2-hydroxy-5-iodobenzoate (70.2 g, 0.22 mol) and 4-ethenyl-N-(3-methyl-2-pyridinyl)benzenesulfonamide (54.8 g, 0.20 mol) were suspended in a solution of tributylamine (55.5 g, 0.3 mol) and palladium(II) acetate (0.44 g 0.0022 mol) in dimethylacetamide (200 ml)

35

The mixture was heated to 100 °C for 3 h. Activated carbon (2 g) was added and the

suspension filtered and the filtrate poured into boiling isopropanol.

The solution was slowly cooled to 10 °C and the crystals were filtered off. The filter cake was washed with cold isopropanol (20 ml) and the product was dried.

5

Yield 78 g, (83 %).

#### Example 12d

10 *2-Hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.*

The product from Example 12c (78 g) was added to a solution of potassium hydroxide (50 g, 0.78 mol) in water (600 ml) and the mixture was refluxed overnight. After addition of activated carbon (2 g) and sodium sulfite (0.5 g) the mixture was allowed to cool, with stirring to room temperature and then filtered.

15

After filtration, ethanol (600 ml) was added and the solution was heated to about 90 °C. Hydrochloric acid (80 ml) was added in one portion and after about 3 min. the product began to precipitate.

20

The suspension was slowly cooled to room temperature and the product was filtered off. The filter-cake was thoroughly washed with water and dried. Yield: 64 g, (78 % over-all).

NMR proved the identity with the product of Example 4.

25

#### Example 13

Isobutyl 2-hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoate

#### Exempel 13a

30 *4-(2-Bromoethyl)-N-(2-pyridinyl)benzenesulfonamide*

This compound was synthesized analogously to Example 12a. Yield 67 %

#### Example 13b

35 *4-Ethenyl-N-(2-pyridinyl)benzenesulfonamide*

This compound was synthesized analogously to Example 12b. Yield 66 %

## Example 13c

*Isobutyl 2-hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoate.*

- Isobutyl 2-hydroxy-5-iodobenzoate (19.2 g 0.06 mol), 4-ethenyl-N-(2-pyridinyl)-benzenesulfonamide (13 g, 0.05 mol) and palladium(II) acetate (0.11 g, 0.0005 mol) were suspended in a mixture of tributylamine (13.8 g, 0.075 mol) and dimethylacetamide (50 ml).

- After heating and stirring for 160 min. at 85 °C acetic acid (5ml) was added and the hot mixture was poured into boiling isopropanol (400 ml). Upon cooling in the refrigerator a precipitate formed which was collected by filtration. The filter-cake was washed with water and isopropanol.

Yield: 4.8 g, (21.2 %).

- NMR proved the identity with the product of Example 5.

## Example 14

2-Hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid acetone solvate.

20

Potassium 2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]-benzoate (28 g) was dissolved in water (160 ml) and acetone (240 ml) with heating to 50 °C. Hydrochloric acid (6 ml) was rapidly added. The product crystallized as a solvate containing about 0.14 mol acetone/mol 2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid. Yield 23.7 g.

25

<sup>1</sup>H-NMR: Spin system A: δ 7.04(d), 7.70(dd), 8.00(d): (=salicylate ring).  
Spin system B: δ 7.70(d,2H), 7.95(d,2H): (=central benzene ring)  
Spin system C: δ 7.66(d, broad)), 6.83(s, broad), 7.84(s, broad): (=pyridine ring).  
Spin system D: δ 2.15(s, broad): (=methyl group)  
Spin system E: δ 2.10(s): (=acetone)

30

- The broadening of signals in the pyridine ring is typical for 3-methyl substituted derivatives. The amount of acetone corresponds to about 0.15 equivalents.

35

## Example 15

2-Hydroxy-5-[2-[4-[(5-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid

## 5 Example 15a

*Tributylammonium 4-ethenyl-benzenesulfonate.*

10 4-(2-Bromoethyl)benzenesulfonyl chloride (89.1 g, 0.36 mol) and potassium hydroxide (86.2 g 1.44 mol) and hydroquinone (0.5 g) were refluxed in water ( 1 l) and ethanol (500 ml) for 3 h.

The solution was concentrated to about 1 l by evaporation in vacuo and acidified with 2M sulfuric acid (about 400 ml). Tributylamine (71.5 g, 0.39 mol) was added and the solution extracted 4 times with dichloromethane (150 ml each). The combined organic phases was washed with 0.5M sulfuric acid (150 ml), dried with  
15 magnesium sulfate and taken to dryness. Yield 134.1 g, quantitative.

## Example 15b

*Methyl 2-hydroxy-5-[2-(4-sulfophenyl)ethenyl]benzoate potassium salt*

20 Tributylammonium 4-ethenyl-benzenesulfonate (134 g, 0.36 mol), methyl 2-hydroxy-5-iodo-benzoate (101 g, 0.36 mol), tributylamine (101 g, 0.54 mol) were dissolved in dimethylacetamide (370 ml). Palladium(II) acetate (0.8 g, 0.0036 mol) was added and the solution was heated with stirring to 85 °C for 3 h. The suspension was filtered to remove insoluble palladium salts and the solution evaporated to  
25 dryness. The residue was dissolved in dichloromethane (500 ml). With vigorous stirring, a solution of potassium hydroxide (94 g, 1.45 mol) in water (300 ml) was quickly added. The precipitated dipotassium salt was filtered off and washed with dichloromethane and diethyl ether. After drying, the solid material was refluxed in acetic acid (200 ml) for 30 min. After cooling to room temperature, the solids were  
30 filtered off and washed with diethyl ether. Yield 57.6 g (43 %).

## Example 15c

*Methyl 2-aceryloxy-5-[2-(4-sulfophenyl)ethenyl]benzoate potassium salt*

35 Methyl 2-hydroxy-5-[2-(4-sulfophenyl)ethenyl]benzoate potassium salt (57.5 g, 0.15 mol), acetic acid (35 ml) and acetic anhydride (142 ml) were refluxed until the major part of the solids had dissolved, and then more acetic anhydride (142 ml) and

sulfuric acid (1 ml) were added. After boiling for 1.5 h, the mixture was cooled to room temperature and diethyl ether was added to complete precipitation. After filtration and drying, the yield was 43.4 g (69 %).

5      Example 15d

*Methyl 2-acetyloxy-5-[2-(4-(chlorosulfonyl)phenyl)ethenyl]benzoate*

Methyl 2-acetyloxy-5-[2-(4-sulfophenyl)ethenyl]benzoate potassium salt (42.4 g, 0.104 mol) was suspended in dichloromethane. Dimethylformamide (5 ml) and  
10      thionyl chloride were added, and the mixture refluxed with stirring for 3 h. After cooling to room temperature, water (15 ml) was added. By careful addition of 5M sodium hydroxide, the pH of the aqueous phase was adjusted to about 7. The phases were separated and the organic phase washed with water. The solution was dried (MgSO<sub>4</sub>), treated with activated carbon and filtered. After evaporation of the  
15      solvent, the solid residue was recrystallized from a toluene and petroleum ethermixture. Yield 29.3 g (71 %).

Example 15e

20      *2-Hydroxy-5-[2-[4-[(5-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid.*

Methyl 2-acetyloxy-5-[2-(4-(chlorosulfonyl)phenyl)ethenyl]benzoate (1.2 g, 0.003 mol) 2-amino-5-methyl-pyridine (0.39 g, 0.0036 mol) were added to dry pyridine (10 ml) and the solution stirred at room temperature over night. The solvent was evaporated and the solid residue stirred with 2M sulfuric acid. After filtration, 1.33 g  
25      intermediate ester was obtained. This was refluxed with sodium hydroxide (0.6 g) in water (20 ml) and ethanol (10 ml) for 5 h. After cooling and acidification with hydrochloric acid, the crystalline precipitate was filtered off and washed with water. After stirring with hot water for 1 h, the material was collected and dried. The material was recrystallized by dissolution dissolution in hot 70 % ethanol with 3  
30      equivalents of sodium hydroxide, followed by acidification by hydrochloric acid. After filtration, washing with water, and drying the yield was 0.7 g, (58 %).

<sup>1</sup>H-NMR:      Spin system A: δ 7.01(d), 7.83(dd), 8.01(d): (=salicylate ring).  
Spin system B: δ 7.72(d,2H), 7.82(d,2H): (=central benzene ring)  
35      Spin system C: δ 7.11(d), 7.56(dd), 7.68(d): (=pyridine ring)  
Spin system D: δ 7.17(d), 7.37(d): (=ethenediyl bridge)  
Spin system E: δ 2.15(s): (=methyl group)

## Example 16

2-Hydroxy-5-[2-[4-[3-(phenylmethoxy)-2-pyridinylamino]sulfonyl]-phenyl]ethenyl]benzoic acid

5 This compound was synthesized analogously to Example 15e. Yield 53 %

10 <sup>1</sup>H-NMR: Spin system A: δ 6.99(d), 7.82(dd), 8.0(d): (=salicylate ring).  
Spin system B: δ 7.71(d,2H), 7.93(d,2H): (=central benzene ring)  
Spin system C: δ 6.9(broad), 7.3(broad), 7.66(d): (=pyridine ring)  
Spin system D: δ 7.17(d), 7.38(d): (=ethenediyl bridge)  
Spin system E: δ 7.48(s), 7.3-7.4(m): (=phenyl group)  
Spin system F: δ 5.15(s): (=methylene group)

## Example 17

15 5-[2-[4-[5-Chloro-2-pyridinylamino]sulfonyl]phenyl]ethenyl]-2hydroxybenzoic acid

This compound was synthesized analogously to Example 15e. Yield 82 %

20 <sup>1</sup>H-NMR: Spin system A: δ 7.02(d), 7.84(dd), 8.04(d): (=salicylate ring).  
Spin system B: δ 7.77(d,2H), 7.91(d,2H): (=central benzene ring)  
Spin system C: δ 7.14(d), 7.81(dd), 8.24(d): (=pyridine ring)  
Spin system D: δ 7.20(d), 7.41(d): (=ethenediyl bridge)

## 25 Example 18

2-Hydroxy-5-[2-[4-[5-methyl-3-isoxazolyl]aminosulfonyl]phenyl]ethenyl]benzoic acid

30 This compound was synthesized analogously to Example 15e. Yield 78 %.

35 <sup>1</sup>H-NMR: Spin system A: δ 7.03(d), 7.85(dd), 8.05(d): (=salicylate ring).  
Spin system B: δ 7.80(d,2H), 7.85(d,2H): (=central benzene ring)  
Spin system C: δ 6.18(s), : (=isoxazole ring)  
Spin system D: δ 7.21(d), 7.43(d): (=ethenediyl bridge)  
Spin system E: δ 2.33(s): (=methyl group).

## Example 19

2-Hydroxy-5-[2-[4-[(4-methyl-2-pyrimidyl)amino-sulfonyl]phenyl]ethenyl]benzoic acid ethanol solvate (2:1)

5 This compound was synthesized analogously to Example 15e. Yield 38 %

10 <sup>1</sup>H-NMR: Spin system A: δ 7.02(d), 7.84(dd), 8.03(d): (=salicylate ring).  
Spin system B: δ 7.76(d,2H), 7.97(d,2H): (=central benzene ring)  
Spin system C: δ 7.20(d), 7.40(dd): (=pyrimidine ring)  
Spin system D: δ 7.20(d), 7.40(d): (=ethenediyl bridge)  
Spin system E: δ 2.30(s): (=methyl group)  
Spin system F: δ 1.05(t, 0.5x3H), 3.42(d, 0.5x2H): (ethanol)

## Example 20

15 2-Hydroxy-5-[2-[4-[(2-pyridazinyl)aminosulfonyl]phenyl]ethenyl]benzoic acid

This compound was synthesized analogously to Example 15e. Yield 67 %

20 <sup>1</sup>H-NMR: Spin system A: δ 7.02(d), 7.84(dd), 8.03(d): (=salicylate ring).  
Spin system B: δ 7.78(d,2H), 7.93(d,2H): (=central benzene ring)  
Spin system C: δ 8.24(s, 2H), 8.40(s): (=pyridazine ring)  
Spin system D: δ 7.20(d), 7.41(d): (=ethenediyl bridge)

## Example 21

25 2-Hydroxy-5-[2-[4-[[6-fluoro-(2-benzo[d]thiazolyl)]aminosulfonyl]-phenyl]ethenyl]benzoic acid

This compound was synthesized analogously to Example 15e. Yield 27 %.

30 <sup>1</sup>H-NMR: Spin system A: δ 7.0(d), 7.85(dd), 8.02(d): (=salicylate ring).  
Spin system B: δ 7.76(d,2H), 7.83(d,2H): (=central benzene ring)  
Spin system C: δ 7.77(dd), 7.25(ddd), 7.30(dd): (=benzothiazole ring)  
Spin system D: δ 7.20(d), 7.38(d): (=ethenediyl bridge)

## Example 22

2-Hydroxy-5-[2-[4-[N-methyl-(2-pyridinyl)aminosulfonyl]phenyl]ethenyl]benzoic acid

5

This compound was synthesized analogously to Example 15e. Yield 81 %.

<sup>1</sup>H-NMR: Spin system A: δ 7.02(d), 7.87(dd), 8.04(d): (=salicylate ring).  
Spin system B: δ 7.56(d,2H), 7.73(d,2H): (=central benzene ring)  
Spin system C: δ 7.59(d), 7.85(ddd), 7.25(dd), 8.32(dd): (=pyridine ring)  
Spin system D: δ 7.20(d), 7.42(d): (=ethenediyl bridge)  
Spin system E: δ 3.2(s): (=methyl group).

10

15

## Example 23

2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid

## Example 23a

*Isobutyl 2-hydroxy-5-[2-(4-sulfophenyl)ethenyl]benzoate dipotassium salt*

20

Tributylammonium 4-ethenyl-benzenesulfonate (55.2 g, 0.15 mol), isobutyl 2-hydroxy-5-iodo-benzoate (47.8 g, 0.36 mol), tributylamine (41.4 g, 0.15 mol) were dissolved in dimethylacetamide (150 ml). Palladium(II) acetate (0.33 g, 0.0015 mol) was added and the solution was heated with stirring to 85 °C for 7 h. After about 3 h a further portion of isobutyl 2-hydroxy-5-iodo-benzoate (23.9 g, 0.18 mol) was added. The reaction mixture was evaporated to dryness. The residue was dissolved in dichloromethane (750 ml). With vigorous stirring, a solution of 0.5M potassium hydroxide (600 ml) was quickly added. The precipitated dipotassium salt was filtered off and washed with dichloromethane. Yield 42 g, (68 %)

25

30

## Example 23b

*Isobutyl 2-acetyloxy-5-[2-(4-sulfophenyl)ethenyl]benzoate potassium salt*

35

Isobutyl 2-hydroxy-5-[2-(4-sulfophenyl)ethenyl]benzoate dipotassium salt (37.2 g, 0.09 mol) was added to a boiling solution of acetic acid (20 ml) and acetic anhydride (80ml). Sulfuric acid (8 ml) was added cautiously and the mixture refluxed until the



major part of the solids had dissolved, and then more acetic anhydride (80 ml) was added. After boiling for 1 h, the mixture was cooled to room temperature. Diethyl ether (350 ml) was added and the suspension filtered and washed with diethyl ether. Yield 36.1 g (88 %)

5

#### Example 23c

##### *Isobutyl 2-acetyloxy-5-[2-(4-(chlorosulfonyl)phenyl)ethenyl]benzoate*

Isobutyl 2-acetyloxy-5-[2-(4-sulfohenyl)ethenyl]benzoate potassium salt (36.1 g, 0.08 mol) was suspended in dichloromethane. Dimethylformamide (20 ml) and thionyl chloride (20 ml) were added, and the mixture refluxed with stirring for 3 h. After cooling to room temperature, water (15 ml) was added. By careful addition of 5M sodium hydroxide, the pH of the aqueous phase was adjusted to about 7. The phases were separated and the organic phase washed with water. The solution was dried (MgSO<sub>4</sub>), treated with activated carbon and filtered. After evaporation of the solvent, the solid residue was recrystallized from a toluene and petroleum ether mixture. Yield 19.5 g (57 %).

20

#### Example 23d

##### *2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]-ethenyl]benzoic acid*

Isobutyl 2-acetyloxy-5-[2-(4-(chlorosulfonyl)phenyl)ethenyl]benzoate (1.99 g, 0.0046 mol) and 2-pyridinamine (1.3 g, 0.0138 mol) was stirred in dichloromethane (10 ml) at 45 °C for 1 h. After cooling to room temperature dichloromethane (20 ml) and 2M sulfuric acid (20 ml) were added. The precipitated solid (1.3 g) was collected and washed with water. The organic phase was taken to dryness and the residue treated with boiling methanol containing ammonia (2 ml) for 20 min. After cooling, 0.6 g was collected which was combined with the first crop. The combined material (1.9 g) was refluxed with potassium hydroxide (1 g) in water (50 ml). After acidification and filtration, the solid residue was dissolved in 70 % ethanol with about 3 equivalents of potassium hydroxide at the boiling point and acidified while boiling with hydrochloric acid. After filtration and drying, the yield was 1.25 g (70 %).

35

NMR proved the identity to the product of Example 3.

## Example 24

2-Hydroxy-5-[2-[4-[(6-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid

5 This compound was synthesized analogously to Example 23d. Yield 69 %

<sup>1</sup>H-NMR: Spin system A:  $\delta$  7.0(d), 7.82(dd), 8.02(d): (=salicylate ring).  
Spin system B:  $\delta$  7.71(d,2H), 7.83(d,2H): (=central benzene ring)  
Spin system C:  $\delta$  7.04(d), 7.61(dd), 6.66(d): (=pyridine ring)  
10 Spin system D:  $\delta$  7.17(d), 7.36(d): (=ethenediyl bridge)  
Spin system E:  $\delta$  2.30(s): (=methyl group).

## Example 25

15 2-Hydroxy-5-[2-[4-[(4-methyl-2-pyridinylamino)sulfonyl]phenyl]-ethenyl]benzoic acid ethanol solvate (2:1)

This compound was synthesized analogously to Example 23d. Yield 69 %.

<sup>1</sup>H-NMR: Spin system A:  $\delta$  7.0(d), 7.83(dd), 8.02(d): (=salicylate ring).  
20 Spin system B:  $\delta$  7.72(d,2H), 7.84(d,2H): (=central benzene ring)  
Spin system C:  $\delta$  7.04(d), 6.69(dd), 7.83(d): (=pyridine ring)  
Spin system D:  $\delta$  7.18(d), 7.37(d): (=ethenediyl bridge)  
Spin system E:  $\delta$  2.25(s): (=methyl group).

## 25 Example 26

2-Hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]benzoic acid

A solution of 2-hydroxy-5-[2-[4-[(2-pyridinylamino)sulfonyl]phenyl]ethenyl]-benzoic acid (3 g, 7.5 mmol) in tetrahydrofuran (100 ml) and acetic acid (100 ml)  
30 was hydrogenated over palladium on charcoal (10 %, 0.3g) and hydrogen at atmospheric pressure and room temperature for 27 h. The catalyst was filtered off and the solvent was evaporated. The residue was dissolved in a solution of sodium hydroxide (0.3g) in acetone (30 ml) and water (30 ml). The solution was acidified with hydrochloric acid at ca 50 °C. After cooling, the product crystallized yielding  
35 1.6g (52 %).

- 5 <sup>1</sup>H-NMR: Spin system A:  $\delta$  6.81(d), 7.29(dd), 7.61(d): (=salicylate ring).  
Spin system B:  $\delta$  7.35(d,2H), 7.76(d,2H): (=central benzene ring)  
Spin system C:  $\delta$  7.13(d), 7.67(ddd), 6.84(ddd), 7.98(d): (=pyridine ring)  
Spin system D:  $\delta$  2.7-2.9(m, 4H): (=ethanediyl bridge)

## Example 27

2-Hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]acetyl]benzoic acid.

- 10 A solution of 2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoic acid (4.1g, 0.01mol) in formic acid (100 ml) was refluxed for 30 h. Water (200 ml) was added and the product crystallized to yield 3.0g (71 %) product.

- 15 <sup>1</sup>H-NMR: Spin system A:  $\delta$  7.08(d), 8.20(dd), 8.48(d): (=salicylate ring).  
Spin system B:  $\delta$  7.42(d,2H), 7.90(d,2H): (=central benzene ring)  
Spin system C:  $\delta$  7.62(d), 6.78(broad), 7.82(broad): (=pyridine ring)  
Spin system D:  $\delta$  4.45(s): (=methylene bridge)  
Spin system E:  $\delta$  2.15(s): (=methyl group).

## 20 Example 28

2-Hydroxy-5-[[4-[(2-pyridinylamino)sulfonyl]phenyl]acetyl]benzoic acid.

The compound was prepared from 2-hydroxy-5-[[4-[(2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoic acid following the method of example 27.

- 25 <sup>1</sup>H-NMR: Spin system A:  $\delta$  7.05(d), 8.14(dd), 8.44(d): (=salicylate ring).  
Spin system B:  $\delta$  7.38(d,2H), 7.81(d,2H): (=central benzene ring)  
Spin system C:  $\delta$  7.16(dd), 7.69(ddd), 6.84(ddd), 7.98(dd): (=pyridine ring)  
30 Spin system D:  $\delta$  4.4(s): (=methylene bridge)

## Example 29

2-Hydroxy-5-[1-oxo-3-[4-[(2-pyridinylamino)sulfonyl]phenyl]-2-propenyl]benzoic acid.

- 35 4-Formyl-N-(2-pyridinyl)benzenesulfonamide (5.7 g, 0.022 mol), 5-acetyl-2-

hydroxybenzoic acid (6.64 g, 0.044 mol) in 5M sodium hydroxide (20 ml) were stirred for 3 days at room temperature. The solution was diluted with water (100 ml) and acetic acid added dropwise in excess. The precipitate was collected by filtration and washed with water. After drying, the material was recrystallized from acetic acid and dried at 120 °C in vacuo. Yield 6.4 g, 66 %.

<sup>1</sup>H-NMR: Spin system A: δ 7.12(d), 8.35(dd), 8.58(d): (=salicylate ring).  
Spin system B: δ 7.93(d,2H), 8.05(d,2H): (=central benzene ring)  
Spin system C: δ 7.23(d), 7.76(ddd), 6.88(ddd), 8.02(m): (=pyridine ring)  
Spin system D: δ 7.77(d), 8.02(d): (=oxopropenediyl bridge)

#### Example 30

2-Hydroxy-5-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl-benzoic acid.

#### Example 30a

*4-Methylphenyl 4-iodobenzenesulfonate.*

4-Iodobenzenesulfonyl chloride (604 g, 2 mol) was added with stirring to a solution of 4-methylphenol (227 g, 2.1 mol) and triethylamine (222 g, 2.2 mol) in toluene (1.5 l). The suspension was stirred at 60 ° C for 2 h. The reaction mixture was added to excess dilute hydrochloric acid. The precipitate was filtered off and washed with ethanol. A second crop was obtained by concentrating the toluene mother liquor and collecting the solid material in the same manner. The combined crops were recrystallized from 2-propanol (2.5 l). Yield 680 g, 91 %.

#### Example 30b

*4-Methylphenyl 4-(2-trimethylsilylethynyl)benzenesulfonate.*

4-Methylphenyl 4-iodobenzenesulfonate (374 g, 1 mol) was dissolved in tetrahydrofuran ( 1 l). Triethylamine (200 g, 2 mol), dichlorobis(triphenylphosphine)palladium (1.1 g, 1.6 mmol) and copper(I)iodide (1.9 g, 10 mmol) were added. The solution was heated to 40 ° C and trimethylsilylethyne (108 g, 1.1 mol) was added during 15 min with stirring and cooling to keep the reaction temperature below 55 ° C. Close to the end of the

addition, the colour of the reaction mixture changed from yellow to brownish green. The reaction mixture was kept at about 40 ° for 30 min, filtered from particulate matter and the solvent evaporated. The residue was dissolved in toluene and washed with dilute hydrochloric acid. After drying and treatment with activated carbon, the solution was evaporated. 392 g of the oily product was obtained, containing some toluene. The material was used directly in the next step without further purification.

#### Example 30c

##### *4-Methylphenyl 4-ethynylbenzenesulfonate.*

The crude oily 4-methylphenyl 4-(2-trimethylsilylethynyl)benzenesulfonate from the preceding step (392 g) was dissolved in dimethylformamide (800 ml). Water (20 ml) and potassium fluoride (50 g, 0.86 mol) were added. The solution turned dark. After 20 min it was diluted with water and the precipitated oily product soon solidified. The material was collected and dissolved in boiling heptane (4 l). The solution was treated with activated carbon and dried with magnesium sulfate. After filtering the solution through a layer (1 cm) of alumina a crystalline product was obtained as three crops by crystallisation and evaporation. The total yield was 242 g, 89% in the two steps from 4-methylphenyl 4-iodobenzenesulfonate.

#### Example 30d

##### *Ethyl 2-hydroxy-5-[[4-(4-methylphenoxy)sulfonyl]phenyl]ethynyl]benzoate.*

Ethyl 2-hydroxy-5-iodobenzoate (58.5 g, 0.2 mol), 4-methylphenyl 4-ethynylbenzenesulfonate (54.4 g, 0.2 mol) and triethylamine (40 g, 0.4 mol) were dissolved in tetrahydrofuran (100 g). The solution was heated to 50 ° C and dichlorobis(triphenylphosphine)palladium (0.5 g, 0.72 mmol) and copper(I)iodide (1 g, 5.2 mmol) were added simultaneously with stirring. The reaction mixture turned dark and triethylamine hydroiodide started to precipitate after about 10 min. After keeping at 45 ° C for 45 min, the suspension was poured into dilute hydrochloric acid and the mixture was extracted with toluene. The toluene solution was evaporated to dryness. The residue was triturated with acetonitrile, filtered off and dried. Yield 73 g, (84 %). The pooled material from several batches (270 g) was recrystallized from acetonitrile (1.2 l). Yield 250 g.

## Example 30e

*Ethyl 2-hydroxy-5-[(4-sulfophenyl)ethynyl]benzoate potassium salt.*

5 Ethanol (500 ml) and toluene (100 ml) was mixed and 200 ml distilled off. Sodium metal (11.5 g, 0.5 mol) was added in portions. The sodium ethoxide solution was heated to boiling and ethyl 2-hydroxy-5-[[4-(4-methylphenoxy)sulfonyl]phenyl]ethynyl]benzoate (43.6 g, 0.1 mol) was added and the solution refluxed for 30 min. The solution was neutralized with acetic acid and cooled. A thick paste was formed. The solid was filtered off and washed with  
10 ethanol. It was dissolved in boiling water and potassium acetate (90 g) was added. The solution was cooled to ambient temperature and the crystalline product filtered off, washed with acetone and dried at 100 ° C. Yield practically quantitative. The product contains some inorganic salts.

## 15 Example 30f

*Ethyl 2-hydroxy-5-[(4-chlorosulfonylphenyl)ethynyl]benzoate.*

Ethyl 2-hydroxy-5-[(4-sulfophenyl)ethynyl]benzoate potassium salt (19 g, 50 mmol) was suspended in toluene (100 ml). Dimethylformamide ( 1 ml) and thionyl chloride  
20 (12 g) were added and the suspension was refluxed for 1 h. The reaction mixture was added to a nearly saturated sodium chloride solution and the precipitated oil was extracted with toluene. The solution was dried and taken to dryness. The product was sufficiently pure for the next step. Yield 16 g, 88 %.

## 25 Example 30g

*Ethyl 2-acetyloxy-5-[(4-chlorosulfonylphenyl)ethynyl]-benzoate.*

Ethyl 2-hydroxy-5-[(4-chlorosulfonylphenyl)ethynyl]benzoate (16 g, 44 mmol) was dissolved in acetic anhydride (40 ml) at 120 ° C. Sulfuric acid (about 0.4 ml) was  
30 added dropwise. After 5 min the solution was diluted with toluene (about 100 ml) and added to saturated sodium chloride solution with vigorous stirring. The phases were separated, the toluene solution dried and evaporated. More toluene was added and the evaporation repeated. The oily product crystallized. The product was sufficiently pure for the next step. Yield 17.0 g (95 %).

35

## Example 30h

*2-Hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]-benzoic acid.*

- 5 3-methyl-2-pyridinamine (1.3 g, 12 mmol) was dissolved in dry pyridine (20 ml) and ethyl 2-acetyloxy-5-[(4-chlorosulfonylphenyl)ethynyl]-benzoate (4.1 g, 10 mmol) was added. The solution was kept at ambient temperature for 18 h. The solvent was evaporated and the residue dissolved in a small amount of tetrahydrofuran. The solution was added to a refluxing solution of potassium hydroxide (6 g) in water (100 ml) and ethanol (50 ml). After 15 min the solution was acidified with formic acid to produce a precipitate. The solid was collected by filtration, washed with water and dried. Yield 1.2 g, 29 %.

NMR proved the identity with the product of Example 2.

15

## Example 31

2-Hydroxy-5-[[4-[[3-(phenylmethoxy)-2-pyridinylamino]sulfonyl]phenyl]ethynyl]benzoic acid.

- 20 This compound was synthesized analogously to Example 30h. The product was recrystallized from acetic acid and then nitromethane. Yield 58 %

- <sup>1</sup>H-NMR: Spin system A: δ 7.06(d), 7.7(dd, overlap with other signals), 8.0(d): (=salicylate ring).  
25 Spin system B: δ 7.7(d, 2H, overlap with other signals), 8.02(d, 2H): (=central benzene ring)  
Spin system C: δ 7.40(d, broad), 6.97(t, broad), 7.7(overlap with other signals): (=pyridine ring)  
Spin system D: δ 7.53(d, 2H), 7.43(t, 2H), 7.36(t): (=phenyl group)  
30 Spin system F: δ 5.10(s): (=methylene group)

## Example 32

5-[[4-[(5-Chloro-2-pyridinylamino)sulfonyl]phenyl]ethynyl]-2hydroxybenzoic acid.

- 35 This compound was synthesized analogously to Example 30h. The product was recrystallized twice from a dimethoxyethane/methylcyclohexane mixture. Yield 58 %

<sup>1</sup>H-NMR: Spin system A: δ 7.04(d), 7.70(dd), 7.99(d): (=salicylate ring).  
Spin system B: δ 7.73(d,2H), 7.94(d,2H): (=central benzene ring)  
Spin system C: δ 7.12(d), 7.82(dd), 8.23(d): (=pyridine ring)

5

## Example 33

2-Hydroxy-5-[[4-[(4-methyl-2-pyridinyl)amino-sulfonyl]phenyl]ethynyl]benzoic acid.

10 This compound was synthesized analogously to Example 30h. The product was dissolved in potassium hydroxide at pH about 9. Potassium acetate was added and the precipitated potassium salt was collected by filtration. The salt was dissolved in water (100 ml) and ethanol (50 ml) and the product precipitated by acidification with formic acid. Hydrochloric acid was then added to decrease the pH to less than 2. The  
15 solid was collected by filtration, washed with water and dried. Yield 2.5 g, 61 %.

<sup>1</sup>H-NMR: Spin system A: δ 7.04(d), 7.7(dd), 7.98(d): (=salicylate ring).  
Spin system B: δ 7.68(d,2H), 7.89(d,2H): (=central benzene ring)  
Spin system C: δ 6.70(dd), 7.07(broad), 7.82(d): (=pyridine ring)  
20 Spin system D: δ 2.24(s): (=methyl group)

## Example 34

2-Hydroxy-5-[[4-[[6-fluoro-(2-benzothiazolyl)aminosulfonyl]phenyl]ethynyl]benzoic acid.

25

This compound was synthesized analogously to Example 30h. The product was dissolved in dilute sodium hydroxide and carefully neutralized to about pH 7. Some solid material was filtered off and the solution reprecipitated by the addition of  
30 formic and hydrochloric acid as before. Yield 2.2g, 47 %.

<sup>1</sup>H-NMR: Spin system A: δ 7.03(d), 7.70(dd), 7.99(d): (=salicylate ring).  
Spin system B: δ 7.74(d,2H), 7.90(d,2H): (=central benzene ring)  
35 Spin system C: δ 7.79(dd), 7.27(ddd), 7.33(dd): (=benzothiazole ring)



## Example 35

2-Hydroxy-5-[[4-[N-methyl-(2-pyridinyl)aminosulfonyl]phenyl]ethynyl]benzoic acid.

5 This compound was synthesized analogously to Example 30h. Yield 2.6 g, 64 %.

<sup>1</sup>H-NMR: Spin system A:  $\delta$  7.05(d), 7.7(dd, overlap with other signals), 8.01(d):  
(=salicylate ring).  
Spin system B:  $\delta$  7.63(d,2H), 7.71(d,2H): (=central benzene ring)  
10 Spin system C:  $\delta$  7.58(ddd), 7.89(ddd), 7.25(ddd), 8.32(ddd):  
(=pyridine ring)  
Spin system E:  $\delta$  3.2(s): (=methyl group).

## 15 Example 36

2-Hydroxy-5-[[4-[(5-methyl-3-isoxazolyl)aminosulfonyl]phenyl]ethynyl]benzoic acid.

## Example 36a

20 *Ethyl 2-hydroxy-5-[[4-[(5-methyl-3-isoxazolyl)aminosulfonyl]phenyl]ethynyl]-benzoate.*

5-methyl-2-isoxazolamine (1.2 g, 12 mmol) was dissolved in dry pyridine (20 ml) and ethyl 2-acetyloxy-5-[(4-chlorosulfonylphenyl)ethynyl]-benzoate (4.1 g, 10  
25 mmol) was added. The solution was kept at ambient temperature for 72 h. The solvent was evaporated and the residue dissolved in a small amount of tetrahydrofuran. The solution was added to a refluxing solution of potassium hydroxide (2.5 g, 38 mmol) in water (100 ml). After 5 min, ethanol (50 ml) was added and the solution acidified with formic acid to produce a precipitate. The solid  
30 was collected by filtration, washed with water and dried. It was recrystallize three times from ethanol. Yield 1.5 g, 35 %.

## Example 36b

35 2-Hydroxy-5-[[4-[(5-methyl-3-isoxazolyl)aminosulfonyl]phenyl]-ethynyl]benzoic acid.

Ethyl 2-hydroxy-5-[[4-[(5-methyl-3-isoxazolyl)aminosulfonyl]-

phenyl]ethynyl]benzoate (1.5 g) was added to a boiling solution of potassium hydroxide (6 g, 92 mmol) in water (100 ml). After 5 min, ethanol (25 ml) was added and the solution acidified with formic acid. The precipitate was collected by filtration, washed with water and dried. Yield 1.4 g, quantitative.

5

<sup>1</sup>H-NMR: Spin system A:  $\delta$  7.02(d), 7.68(dd), 7.97(d): (=salicylate ring).  
Spin system B:  $\delta$  7.74(d,2H), 7.87(d,2H): (=central benzene ring)  
Spin system C:  $\delta$  6.14(s), : (=isoxazole ring)  
Spin system E:  $\delta$  2.25(s): (=methyl group).

10

#### Example 37

4-Fluoro-2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]-benzoic acid.

15

#### Example 37a

*Ethyl 4-fluoro-2-hydroxybenzoate.*

20

4-Fluoro-2-hydroxy-benzoic acid (10.8 g, 0.07 mol), sodium acetate (6.9 g, 0.084 mol), ethyl iodide (55 g, 0.35 mol) and dimethylformamide was stirred at ambient temperature for 48 h. The mixture was poured into water and extracted with n-hexane. The solution was filtered through a plug of silica gel and evaporated. The residue was almost pure product and was used directly in the next step. Yield 10.2 g, 80 %.

25

#### Example 37b

*Ethyl 4-fluoro-2-hydroxy-5-iodobenzoate.*

30

Ethyl 4-fluoro-2-hydroxybenzoate (10.2 g, 55 mmol) was dissolved in n-heptane. Iodo monochloride (12 g, 74 mmol) was added dropwise at ambient temperature with stirring. After 1 h, water was added and then solid sodium sulfite in small portions until the suspension was decolourized. More n-heptane was added to clear solution and the phases separated. The organic phase was dried and concentrated. The product crystallized on cooling. It was collected by filtration and washed with cold n-heptane. The crude yield was 10 g. The product was a mixture of ethyl 4-fluoro-2-hydroxy-5-iodobenzoate (about 80 %) and the 5 chloro analogue (about 20 %).

35

## Example 37c

*Ethyl 4-fluoro-2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]-phenyl]ethynyl]benzoate.*

5

Ethyl 4-fluoro-2-hydroxy-5-iodobenzoate (3.6 g, 80%, 10 mmol), 4-ethynyl-N-(3-methyl-2-pyridinyl)benzenesulfonamide (2.7 g, 10 mmol) and triethylamine (5 g, 50 mmol) were suspended in tetrahydrofuran (20 ml) and the mixture heated to 50 ° C. Dichlorobis(triphenylphosphine)palladium (45 mg, 0.06 mmol) and copper(I)iodide (76 mg, 0.4 mmol) were added with stirring. After about 15 min a practically clear solution had formed. After 3 h at 50 ° C the solution was taken to dryness and triturated with isobutylmethylketone and dilute hydrochloric acid. The organic phase was separated and the solvent evaporated. The oily residue was triturated with a small amount of acetonitrile to form crystals. These were recrystallized from acetic acid. The product was dissolved in a small amount of chloroform and the solution applied on top of a short silica gel column and eluted with 25% isobutylmethylketone in toluene. The pure product resulted after evaporation of the solvent. Yield 2.0 g, 44 %.

20

## Example 37d

*4-Fluoro-2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]-benzoic acid.*

25

Ethyl 4-fluoro-2-hydroxy-5-[[4-[(3-methyl-2-pyridinylamino)-sulfonyl]-phenyl]ethynyl]benzoate (2.0 g, 4.4 mmol) was added to a boiling solution of potassium hydroxide (2.0 g) in water (50 ml). After 15 min water (50 ml) and ethanol(50 ml) were added and the solution acidified with formic acid and hydrochloric acid. After cooling the precipitate was filtered off and dried. Yield quantitative.

30

<sup>1</sup>H-NMR. (Solvent: trifluoroacetic acid)

Spin system A: δ 6.80(d), 8.21(d): (=salicylate ring).

Spin system B: δ 7.72(d,2H), 7.91(d,2H): (=central benzene ring)

Spin system C: δ 7.64(dd), 8.31(d), 8.45(d): (=pyridine ring)

35

Spin system D: δ 2.35(s): (=methyl group)

## Example 38

4-Fluoro-2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethenyl]-benzoic acid.

## 5 Example 38a

*Ethyl 4-fluoro-2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)-sulfonyl]-phenyl]ethenyl]benzoate.*

10 4-Ethenyl-N-(3-methyl-2-pyridinyl)benzenesulfonamide (2.7 g, 10 mmol), ethyl 4-fluoro-2-hydroxy-5-iodobenzoate (3.6 g, 80%, 10 mmol), triethylamine (5 ml) and N,N-dimethylacetamide (5 ml) were mixed and heated to 50° C. Palladium acetate (40 mg) was added and the solution kept at 50° C for 18 h. The solution was poured into water and the precipitate dissolved in chloroform. The solution was dried, applied on top of a short silica gel column and eluted with 30%  
15 isobutylmethylketone in toluene. The pure fractions were collected and evaporated to dryness. After trituration with acetonitril the product crystallized. Yield 1.1 g.

## Example 38b

20 *4-Fluoro-2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)-sulfonyl]-phenyl]ethenyl]benzoic acid.*

25 Ethyl 4-fluoro-2-hydroxy-5-[2-[4-[(3-methyl-2-pyridinylamino)-sulfonyl]-phenyl]ethenyl]benzoate (1.1 g, 2.4 mmol) was added to a boiling solution of potassium hydroxide (2.0 g) in water (50 ml). Ethanol (10 ml) was added to produce a clear solution. After 15 min ethanol (25 ml) were added and the solution neutralized to pH about 7 with formic acid. The slightly opalescent solution was treated with decolourizing carbon and acidified with formic acid. After cooling the precipitate was filtered off and dried. Yield 0.8 g

30 <sup>1</sup>H-NMR. (Solvent: trifluoroacetic acid)

Spin system A: δ 6.79(d), 8.29(d): (=salicylate ring).

Spin system B: δ 7.72(d,2H), 7.90(d,2H): (=central benzene ring)

Spin system C: δ 7.62(dd), 8.29(d), 8.44(d): (=pyridine ring)

Spin system D: δ 7.21(d), 7.35(d): (=ethenediyl bridge)

35 Spin system E: δ 2.36(s): (=methyl group)

## Example 39

2-Hydroxy-5-[[3-[(3-methyl-2-pyridinylamino)sulfonyl]phenyl]ethynyl]benzoic acid.

## 5 Example 39a

*3-Iodo-N-(3-methyl-2-pyridinyl)benzenesulfonamide.*

32.4 g 3-methyl-2-pyridinamine (32.4 g, 0.3 mol) was dissolved in isobutylmethylketone (200 ml) and 3-iodo-benzenesulfonyl chloride (30.2 g, 0.1  
10 mol) was added with stirring. The solution was kept at 40 ° C for 8 h and then added to dilute hydrochloric acid. The precipitate was filtered off and the organic phase was separated and evaporated. The residue was combined with the first solid material and recrystallized from ethanol. The crystals were dissolved in 1M sodium hydroxide (about 200? ml) and extracted with isobutylmethylketone (3 x 100 ml).  
15 The aqueous phase was partially evaporated to remove isobutylmethylketone residues. Ethanol (100? ml) was added and the solution acidified with acetic acid. The precipitate was filtered off, washed with water and dried. Yield 23 g, 61 %.

## Example 39b

20 *Ethyl 2-hydroxy-5-[[3-[(3-methyl-2-pyridinylamino)sulfonyl]-phenyl]ethynyl]-benzoate.*

3-Iodo-N-(3-methyl-2-pyridinyl)benzenesulfonamide (18.7 g, 50 mmol), ethyl 5-ethynyl-2-hydroxybenzoate (9.5 g, 50 mmol), triethylamine (65 ml) and  
25 tetrahydrofuran (65 ml) were stirred at 60 ° C. Dichlorobis(triphenylphosphine)-palladium (250 mg, 0.36 mmol) and copper(I)iodide (140 mg, 0.7 mmol) were added with stirring. After 1 h the solution was poured into dilute hydrochloric acid and extracted with ethyl acetate. The organic phase was washed with water, dried, treated with activated carbon and the solvent evaporated. The residue was recrystallized  
30 from methanol, then from toluene and finally from acetic acid. Yield 12.7 g 58 %.

## Example 39c

*2-Hydroxy-5-[[3-[(3-methyl-2-pyridinylamino)sulfonyl]-phenyl]-ethynyl]benzoic acid.*

35

Ethyl 2-hydroxy-5-[[3-[(3-methyl-2-pyridinylamino)sulfonyl]-phenyl]-ethynyl]benzoate (4 g, 9.2 mmol) was added to a boiling solution of potassium

hydroxide (6 g, 90 mmol) in water (100 ml). After 15 min ethanol (50 ml) was added and the solution acidified with hydrochloric acid. the product was collected by filtration and dried at 110 ° C. Yield 3.5 g, 93 %.

- 5     <sup>1</sup>H-NMR:     Spin system A: δ 7.0(d), 7.68(dd), 7.97(d): (=salicylate ring).  
                         Spin system B: δ 7.55(t), 7.69(d), 7.90(d), 8.06(s): (=central benzene ring)  
                         Spin system C: δ 7.62(d, broad), 6.78(t, broad), 7.81(s, broad): (=pyridine ring)
- 10                   Spin system D: δ 2.10(s): (=methyl group).

The broadening of signals in the pyridine ring is typical for 3-methyl substituted derivatives.

**Example 40****Effect of compounds on human lymphocyte proliferation**

5 Peripheral blood mononuclear cells were isolated from heparinized blood from healthy volunteers by density gradient centrifugation as described by Bain and Pshyk, Transplantation Proc, 4: 163-164 (1972). Cell proliferation induced by Concanavalin A was measured in a standard assay such as described in Söderström  
10 et al., Scand J Immunol, 32:503-516 (1990) using 5% fetal bovine serum in the culture medium. The reaction mixture (final concentrations) consisted of 200 000 cells stimulated with 2.5 µg/ml Concanavalin A and test compounds in a total volume of 0.2 ml. The test compounds were added at the start of the assay, which was run for a total of three days with addition of radioactive [3H]-thymidine during  
15 the last 18 h. Cell bound radioactivity was measured in a liquid scintillation counter.

Table

5 Inhibitory effects of 250  $\mu$ M test compounds on Concanavalin A-induced proliferation of human lymphocytes. Data are from 2 experiments given as per cent inhibition of proliferation in the absence of test compounds.

10	Test compound From Example	Inhibition, %	
		Mean	S.D.
15	Sulfasalazine	53.7	5.2
	1	95.6	2.9
	2	99.5	0.2
	3	73.6	21.2
	4	92.3	4.6
20	15	93.9	2.1
	16	95.4	0.7
	17	99.5	0.2
	18	43.8	26.9
	19	10.3	11.9
25	20	34.8	10.2
	21	99.8	0.2
	22	99.8	0.1
	26	16.0	11.5
	27	37.3	23.4
30	28	15.0	13.1
	29	66.0	4.7
	31	100	0
	32	100	0
	34	98	3
35	35	99	2
	37	100	0
	38	100	0
	39	100	0



**Example 41****Effects of test compounds on superoxide production of human granulocytes.**

- 5 Granulocytes were isolated from the heparinized blood from healthy volunteers. The superoxide assay was essentially that of JM McCord and I Fridovich (J. Biol. Chem. 244:6049-6055 (1969)) as modified by G Carlin et al. (Pharmacol. Toxicol 65:121-127 (1989)). The reaction mixture contained (final concentrations) 125  $\mu$ M cytochrome C in Dulbeccos phosphate buffer with Mg and Ca, 10  $\mu$ M test
- 10 compound and 400 000 granulocytes treated with 5  $\mu$ g/ml of cytochalasin B immediately before the experiment. This reaction mixture was preincubated at 37 °C for 10 minutes, whereafter 10 nM N-formyl-L-methionyl-L-leucyl-L-phenylalanine was added to start the reaction, making the final volume 1 ml. After 10 minutes the tubes were centrifuged and the absorbance of the supernatant at 540 and 550 nm was
- 15 measured. The superoxide production was expressed as the difference in absorbance at these wavelengths.

Table

Inhibitory effects of 10  $\mu$ M test compound on FMLP-induced superoxide production.  
Data are given as per cent inhibition of superoxide production in the absence of test  
5 compounds.

	Test compound From Example	Inhibition, %		Number of exp.
		Mean	S.D.	
10	Sulfasalazine	63.5	11.8	19
	1	71.0	13.6	4
	2	86.0	7.3	4
	3	55.7	17.0	4
15	4	76.9	10.6	4
	5	25.8	10.1	3
	16	94.6	1.1	3
	17	71.9	5.9	3
	18	44.2	6.4	3
20	19	65.9	2.9	3
	20	35.1	7.9	3
	21	93.0	4.2	3
	22	85.5	8.2	4
	25	45.5	4.1	3
25	26	26.8	16.2	4
	27	32.7	17.1	4
	28	17.9	8.7	4
	29	33.4	11.9	5

## Example 42

5 Effects of test compounds on superoxide production of human granulocytes at 100  $\mu$ M concentration.

Granulocytes were isolated from the heparinized blood from healthy volunteers. The superoxide assay was essentially that of JM McCord and I Fridovich (J. Biol. Chem. 244:6049-6055 (1969)) as modified by G Carlin et al. (Pharmacol. Toxicol 65:121-127 (1989)). The reaction mixture contained (final concentrations) 125  $\mu$ M cytochrome C in Dulbeccos phosphate buffer with Mg and Ca, 100  $\mu$ M test compound and 600 000 granulocytes treated with 5  $\mu$ g/ml of cytochalasin B immediately before the experiment. This reaction mixture was preincubated at 37 °C for 10 minutes, whereafter 100 nM N-formyl-L-methionyl-L-leucyl-L-phenylalanine was added to start the reaction, making the final volume 1 ml. After 10 minutes the tubes were centrifuged and the absorbance of the supernatant at 540 and 550 nm was measured. The superoxide production was expressed as the difference in absorbance at these wavelengths.

## 20 Table

Inhibitory effects of 100  $\mu$ M test compound on FMLP-induced (100 nM) superoxide production. Data are given as per cent inhibition of superoxide production in the absence of test compounds.

25

Test compound From Example	Inhibition, %		Number of exp.
	Mean	S.D.	
Sulfasalazine	65.4	9.0	3
2	94.1	4.2	3
33	92.1	3.7	3
37	97.5	3.2	3
38	98.7	1.7	3
39	93.5	1.8	3


35

CLAIMS

1. A compound of the formula  $\text{Het-NR-SO}_2\text{-Ph}^1\text{-A-Ph}^2(\text{COOH})(\text{OH})$  and tautomeric forms, salt, solvate,  $\text{C}_{1-6}$ -alkyl ester and pharmaceutical composition of the compound, in which compound  $\text{Ph}^1$  and  $\text{Ph}^2$  are optionally substituted benzene rings with the condition that carboxy and hydroxy are ortho to one another, A is a bridge and Het- includes an optionally substituted heterocyclic ring which includes conjugated double bonds and binds to NR, **characterized** in that A is a bridge which is stable against reduction because it is not azo; and in that R is hydrogen or lower alkyl.
2. The compound according to Claim 1, **characterized** in that A and  $\text{SO}_2$  bind para or meta to one another on  $\text{Ph}^1$  and that hydroxy and A bind para to one another on  $\text{Ph}^2$ .
3. The compound according to any one of Claims 1-2, **characterized** in that -A- is a straight carbon chain having at most three carbon atoms including one C-C-, C=C- or C $\equiv$ C-bond, optionally together with a carbonyl (C=O).
4. The compound according to any one of Claims 1-3, **characterized** in that -A- is chosen from among -C $\equiv$ C-, -CH=CH-, -CH<sub>2</sub>-CH<sub>2</sub>-, -CO-CH=CH-, -CH=CH-CO- and -CH<sub>2</sub>-CO-, when applicable preferably in the trans-form.
5. The compound according to any one of Claims 1-4, **characterized** in that -A- is chosen from among -C $\equiv$ C- and -CH=CH-.
6. The compound according to any one of Claims 1-5, **characterized** in that R is hydrogen.
7. The compound according to any one of Claims 1-5, **characterized** in that R is lower alkyl, preferably methyl.
8. The compound according to any one of Claims 1-7, **characterized** in that the heterocyclic ring is five-membered or six-

membered with two and three conjugated double bonds respectively.

9. The compound according to Claim 8, **characterized** in that Het is



5 where Het' is the heterocyclic ring and satisfies the formula:



where the free valency binds to NR, and X is:

- 10 (i) -N=CH-NH-, -N=CH-S-, -N=CH-O-, -NH-N=CH-, -O-CH=CH-,  
-CH=CH-O-, -NH-CH=CH-, -CH=CH-NH-, -CH=CH-S-,  
-CH=N-NH-,  
(ii) -CH=CH-CH=CH-, -CH=CH-CH=N-, -CH=N-CH=CH-,  
-CH=CH-N=CH-, -N=CH-CH=CH-,

15 where mutually adjacent bold-typed hydrogen atoms (H) can be substituted in pairs with -CH=CH-CH=CH-, so as to form a bicyclic structure (condensed rings);  
and

$R_1$ ,  $R_2$  and  $R_3$  are substituents on carbon atoms in Het' and are chosen from among the group hydrogen, lower alkyl, halogen, hydroxy, cyano, carboxy, lower alkoxy, benzyloxy, lower acyl (including acetyl and benzoyl) and phenyl and benzyl,  
20 wherein any occurrent benzene rings can be substituted.

10. The compound according to Claim 9, **characterized** in that X is:

-O-CH=CH-, -N-CH=CH-, -CH=CH-NH-, and -CH=CH-S-

25 11. The compound according to Claim 9, **characterized** in that X is:

-CH=CH-CH=CH-, -CH=N-CH-CH-, -CH=CH-N=CH- and  
-N=CH-CH=CH-.

12. The compound according to any one of Claims 9-11,

30 **characterized** in that Het' is a monocyclic structure.

13. The compound according to any one of Claims 9-11,

**characterized** in that Het' is a bicyclic structure.

35 14. The compound according to any one of Claims 9-13,

**characterized** in that  $R_1$  and  $R_2$  are hydrogen and  $R_3$  is chosen from among remaining substituents.

15. The compound according to Claim 14, **characterized** in that  $R_3$  is hydrogen.
- 5 16. The compound according to any one of Claims 12 and 14-15, **characterized** in that Het' is pyridine.
17. The compound according to Claim 16, **characterized** in that A is  $-C\equiv C-$  or  $-CH=CH-$ , and in that  $R_3$  is not hydrogen, preferably methyl.
- 10 18. The compound according to Claim 17, **characterized** in that R is hydrogen.
- 15 19. The compound according to Claim 17, **characterized** in that  $R_3$  is ortho to the  $-NRSO_2-$  substituent in the pyridine ring.
- 20 20. The preparation of a compound according to any one of Claims 1-19, **characterized** in that
- (I) a compound of the formula  $Het-NR-SO_2-Ph^1-X$  is reacted chemically with a compound of the formula  $Y-Ph^2(COOR_5)(OH)$  to coupling between X and Y so as to form a compound according to any one of Claims 1-16, wherein Het, R,  $Ph^1$ ,  $Ph^2$  and A have the same
- 25 significance as in Claim 1;  $R_5$  is hydrogen or lower alkyl; and -X and Y- are pairwise  $-C\equiv CH$  and  $Z_1-$ ;  $-Z_1$  and  $HC\equiv C-$ ;  $-CH=CH_2$  and  $Z_1-$ ;  $-Z_1$  and  $CH_2=CH-$ ;  $-CO-CH_3$  and  $HCO-$ ;  $-CHO$  and  $CH_3-CO-$ ;  $-CH_2COZ_2$  and  $H-$ ;  $-CHO$  and  $Z_3CH_2-$  and  $-CH_2Z_3$  and  $HCO-$ , wherein  $Z_1$  is bromine or iodine,  $Z_2$  is halogen, preferably chlorine, and
- 30  $Z_3$  is the residue of a phosphorous organic compound of Wittig type, for instance triphenylphosphonium group or dialkyl phosphate group, whereafter, when  $R_5$  is not hydrogen, the resultant compound, if so desired, is subjected to a base-catalyzed hydrolysis process to replace said group  $R_5$  with a hydrogen atom or a suitable metal ion; or
- 35 (II) a compound according to formula  $Het-NRH$  is reacted chemically in the presence of a basic condensation agent in a solvent with a lower alkyl ester of a compound according to formula

5 CISO<sub>2</sub>-Ph<sup>1</sup>-A-Ph<sup>2</sup>(COOH)(OR<sub>6</sub>), in which compounds Het, R, Ph<sup>1</sup>, Ph<sup>2</sup> and A have the same significance as in Claim 1 and R<sub>6</sub> is lower acyl, preferably acetyl, so that coupling takes place at the groups -NRH and CISO<sub>2</sub>- to form a compound which with subsequent hydrolysis of occurrent ester groups provides a compound according to any one of Claims 1-16;

10 (III) the bridge A' in a compound of the formula  
 Het-NR-SO<sub>2</sub>-Ph<sup>1</sup>-A'-Ph<sup>2</sup>(COOR<sub>7</sub>)(OR<sub>8</sub>), where Het, R, Ph<sup>1</sup> and Ph<sup>2</sup> have the same significance as in Claim 1; R<sub>7</sub> is hydrogen or R<sub>5</sub>; R<sub>8</sub> is hydrogen or R<sub>6</sub>; and A' can be transformed via an addition reaction or an elimination reaction, alternatively via an oxidation reaction or reduction reaction to A in accordance with Claim 1, is chemically  
 15 transformed to A via one of the aforesaid reactions, whereafter when R<sub>7-8</sub> is separate from hydrogen, the resultant compound, if so desired, is subjected to base catalyzed hydrolysis so as to replace said group R<sub>7-8</sub> with a hydrogen atom or a suitable metal ion.

20 21. A method of treating an autoimmune disease,  
 characterized by administering a therapeutically active dosage of a substance according to Claims 1-19 to an individual suffering from said disease.

22. A compound according to any one of Claims 1-19 for use as a drug.

25 23. The use of a compound according to any one of Claims 1-19 in the preparation of a pharmaceutical composition for treating an autoimmune disease.

# INTERNATIONAL SEARCH REPORT

International Application No PCT/SE 92/00758

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC5: C 07 D 213/75, 261/16, 239/42, 237/20, 277/82											
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched<sup>7</sup></div> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%; border-bottom: 1px solid black;">Classification System</td> <td style="border-bottom: 1px solid black;">Classification Symbols</td> </tr> <tr> <td style="height: 40px; vertical-align: bottom;">IPC5</td> <td style="height: 40px; vertical-align: bottom;">C 07 D</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in Fields Searched<sup>8</sup></div> <p>SE,DK,FI,NO classes as above</p>			Classification System	Classification Symbols	IPC5	C 07 D					
Classification System	Classification Symbols										
IPC5	C 07 D										
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%;">Category *</th> <th style="width: 60%;">Citation of Document,<sup>11</sup> with indication, where appropriate, of the relevant passages<sup>12</sup></th> <th style="width: 30%;">Relevant to Claim No.<sup>13</sup></th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top;">A</td> <td>US, A, 2396145 (ERIC EMIL ANDERS ASKELOF ET AL.) 5 March 1946, see the whole document ---</td> <td style="text-align: center; vertical-align: top;">1-20,22- 23</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">A</td> <td>US, A, 3915951 (KARL HUBERT AGBACK ET AL.) 28 October 1975, see the whole document -----</td> <td style="text-align: center; vertical-align: top;">1-20,22- 23</td> </tr> </tbody> </table>			Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>	A	US, A, 2396145 (ERIC EMIL ANDERS ASKELOF ET AL.) 5 March 1946, see the whole document ---	1-20,22- 23	A	US, A, 3915951 (KARL HUBERT AGBACK ET AL.) 28 October 1975, see the whole document -----	1-20,22- 23
Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>									
A	US, A, 2396145 (ERIC EMIL ANDERS ASKELOF ET AL.) 5 March 1946, see the whole document ---	1-20,22- 23									
A	US, A, 3915951 (KARL HUBERT AGBACK ET AL.) 28 October 1975, see the whole document -----	1-20,22- 23									
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>* Special categories of cited documents:<sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>											
<b>IV. CERTIFICATION</b> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black;">Date of the Actual Completion of the International Search</td> <td style="width: 50%; border-bottom: 1px solid black;">Date of Mailing of this International Search Report</td> </tr> <tr> <td style="height: 40px; vertical-align: bottom;">8th February 1993</td> <td style="height: 40px; vertical-align: bottom; text-align: center;">19 02 1993</td> </tr> <tr> <td style="border-bottom: 1px solid black;">International Searching Authority</td> <td style="border-bottom: 1px solid black;">Signature of Authorized Officer</td> </tr> <tr> <td style="text-align: center; vertical-align: bottom;">SWEDISH PATENT OFFICE</td> <td style="text-align: center; vertical-align: bottom;">Göran Karlsson</td> </tr> </table>			Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	8th February 1993	19 02 1993	International Searching Authority	Signature of Authorized Officer	SWEDISH PATENT OFFICE	Göran Karlsson	
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report										
8th February 1993	19 02 1993										
International Searching Authority	Signature of Authorized Officer										
SWEDISH PATENT OFFICE	Göran Karlsson										

Form PCT/ISA/210 (second sheet) (January 1985)



## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE<sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 21, because ~~they relate to~~ <sup>it is</sup> subject matter not required to be searched by this Authority, namely:

A method for treatment of the human or animal body by therapy,  
see rule 39.1.

2. ☒ Claim numbers 1-9 20 and 22-23, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

These claims are too broadly formulated to permit a meaningful search.  
The search has thus been limited to the compounds considered to be most relevant.

3. ☐ Claim numbers                     , because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING<sup>2</sup>

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the the claims. It is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

## Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO. PCT/SE 92/00758

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the Swedish Patent Office EDP file on 08/01/93  
The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 2396145	46-03-05	NONE	
US-A- 3915951	75-10-28	AT-B- 321305	75-03-25
		AT-B- 329057	76-04-26
		AU-B- 464869	75-09-11
		AU-D- 4917772	74-05-23
		BE-A- 791889	73-05-24
		CA-A- 971954	75-07-29
		CH-A- 553782	74-09-13
		CH-A- 573913	76-03-31
		DE-A-C- 2257629	73-05-30
		FR-A-B- 2161104	73-07-06
		GB-A- 1388920	75-03-26
		JP-A- 48057978	73-08-14
		NL-A- 7215980	73-05-29
		SE-B-C- 374375	75-03-03